The proof of the insolubility in natural numbers for $n > 2$, the Fermat's Last Theorem and Beal's conjecture for coprime integers arranged in a pair A, B, D in the equations

$$A^n + B^n = D^n \text{ and } A^n + B^y = D^z. \\ \text{(elementary aspect)}$$

PROF. DR. K. RAJA RAMA GANDHI¹ AND REUVEN TINT²

Resource person in Math for Oxford University Press and Professor at BITS-Vizag¹

Number Theorist, Israel²

Email: editor126@gmail.com, reuven.tint@gmail.com

Abstract. We give the corresponding identities for different solutions of the equations:

$$aA^x + bB^x = cD^x \text{ [1] and}$$

$$aA^x + bB^y = cD^z \text{ [2]:}$$

As for coprime integers a, b, c, A, B, D and arbitrary positive integers x, y, z further, for not coprime integers, if

$$A_0^{x_0} + B_0^{x_0} = D_0^{x_0} \text{ [3] and}$$

$$A_0^{x_0} + B_0^{y_0} = D_0^{z_0} \text{ [4], where}$$

$x_0, y_0, z_0, A_0, B_0, D_0$ - are any solutions in positive integers.

Theorem.

1. "If [3] and [4] have any solutions $x_0, y_0, z_0, A_0, B_0, D_0$ - in positive integers, then from [1] and [2] we obtain an infinite number of solutions for arbitrary x, y, z ".

Proof.

1.1. Obviously, with respect to

$$a = kD_0^{x_0} + b; c = kA_0^{x_0} + b \text{ and}$$

$$a = kD_0^{z_0} + b; c = kA_0^{x_0} + b, \text{ where}$$

" k " and " b " – are arbitrary positive integers, then

$$(kD_0^{x_0} + b)A_0^{x_0} + bB_0^{x_0} \equiv (kA_0^{x_0} + b)D_0^{x_0} \text{ [5],}$$

$$(kD_0^{z_0} + b)A_0^{x_0} + bB_0^{y_0} \equiv (kA_0^{x_0} + b)D_0^{z_0} [6]$$

1.2. For [3] there exists $k = k'A_0^x B_0^x D_0^x$ and $b = b'A_0^x B_0^x D_0^x$,
where $x; (k', b') = 1$ –are arbitrary positive integers, since [5] we have

$$\begin{aligned} B_0^x D_0^x (k'D_0^{x_0} + b')A_0^{x+x_0} + A_0^x D_0^x B_0^{x+x_0} &\equiv \\ &\equiv A_0^x B_0^x (k'A_0^{x_0} + b')D_0^{x+x_0} [7]. \end{aligned}$$

1.3. Similarly, 1.2. from [6] it follows that

$$B_0^y D_0^z (k'D_0^{z_0} + b')A_0^{x+x_0} + A_0^x D_0^z B_0^{x+y_0} \equiv A_0^x B_0^y (k'A_0^{x_0} + b')D_0^{x+z_0} [8].$$

1.4. If A_0, B_0, D_0 –are for coprime integers arranged in a pair, then equations [5], [6], [7], [8] have an infinite set of solutions for each $n = x + x_0$, this means, that using Faltings (4) the equations [3] and [4] for each $n = x + x_0 > 2$ does not have solutions in natural numbers. Then, $x = 0$ and $x_0 \leq 2$. This means that Fermat's Last Theorem and Beal's conjecture based on previous articles completely resolved. The proof is completed. In the same way [7], [8] instead of x, x_0 holds y, y_0 or z, z_0

Examples:

1.4.1.

$$13^2 \times 3^4 + 11^2 \times 2^4 = 5^2 \times 5^4$$

$$3^4 \times 2^4 \neq 5^4, \text{ but}$$

$$39^2 \times 3^2 + 11^2 \times 4^2 = 5^4 \times 5^2$$

$$39^2 = 56 \times 5^2 + 11^2$$

$$5^4 = 56 \times 3^2 + 11^2$$

$$7^2 + 2^5 = 3^4$$

1.4.2. Let

$$A^2 + B^2 = D^2$$

then, if A and B are not coprime, but

$$3^2 + 4^2 = 5^2,$$

therefore from [7]

$$\begin{aligned} 4^x \times 5^x (k'5^2 + 3^x b') 3^{x+2} + 3^x 5^x 4^{x+2} &\equiv \\ &\equiv 3^x \times 4^x \times (k' \times 3^x + 5^x \times b') \times 5^{x+2} \text{ etc.} \end{aligned}$$

1.4.3. If

$$A_0^n + B_0^n = D_0^n, \text{ then}$$

$$a^2 A_0^n + b^2 B_0^n \equiv c^2 D_0^n \text{ and } a = p^2 D_0^n - q^2 B_0^n,$$

$$b = p^2 D_0^n - 2pqD_0^n + q^2 B_0^n, \text{ where}$$

p, q — are arbitrary natural positive integers.

If $n = 2, p = 2, q = 1; D_0 = 3; B_0 = 4, D_0 = 5$, then

$$7^2 \times 3^4 + 4^2 \times 2^4 = 13^2 \times 5^2,$$

$$21^2 \times 3^2 + 4^2 \times 4^2 = 13^2 \times 5^2,$$

$$209^2 \times 3^2 + 91^2 \times 4^2 = 145^2 \times 5^2 (p = 3; q = 1) \text{ etc.}$$

1.4.4. If

$$a + b = c,$$

then

$$a(p^2 c - q^2 b)^2 + b(p^2 c - 2pqc + q^2 b)^2 \equiv$$

$$\equiv c(p^2 c - 2pqb + q^2 b)^2, \text{ where}$$

p, q — are arbitrary positive integers.

In addition, suppose that:

$$a' = a^n; b' = b^n; c' = c^n;$$

$$a' = a^n; b' = b^y; c' = c^z.$$

References:

- [1] PROF. DR. K. RAJA RAMA GANDHI, Reuven Tint “ The reproductive solution for Fermat’s Last theorem (elementary aspect)- first proof.
- [2] Reuven Tint, The Identities of Ordinary Which Is Leading to the Extraordinary Consequences (Elementary Aspect) Asian Journal of Mathematics and Applications - ISSN 2307-7743
- [3] PROF. DR. K. RAJA RAMA GANDHI AND REUVEN TINT Proof of Beal's Conjecture.
- [4] Faltings G (1983). "Endlichkeitssätze für abelsche Varietäten über Zahlkörpern". *Inventiones Mathematicae* 73 (3): 349–366. doi:10.1007/BF01388432.
- [5] H.Davenport, “The Higher Arithmetic”, Moscow, 1965 [3], p157-163, comments VII p.171).