

# A PROOF OF THE VERACITY OF BOTH GOLDBACH AND DE POLIGNAC CONJECTURES

Jamel Ghanouchi

6 Rue Khansa 2070 Marsa Tunisie

**Keywords** : Goldbach ; De Polignac ; Twin primes ; Algebraic proof

## Abstract

The present algebraic development begins by an exposition of the data of the problem. The definition of the primal radius  $r > 0$  is : For all positive integer  $x \geq 3$  exists a finite number of integers called the primal radius  $r > 0$ , for which  $x + r$  and  $x - r$  are prime numbers. The contrary is that  $2x = (x + r) + (x - r)$  is always the sum of a finite number of primes. Also, for all positive integer  $x \geq 0$ , exists an infinity of integers  $r > 0$ , for which  $x + r$  and  $r - x$  are prime numbers. The conclusion is that  $2x = (x + r) - (r - x)$  is always an infinity of differences of primes.

## Introduction

There is a similarity between the assertion : “an even number is always the sum of two primes” and the assertion : “an even number is always the difference of two primes”. The present article gives the proof that the two assertions are the consequences of the same concept by the introduction of the notion of the primal radius.

## The proof

Let us suppose that exists an integer  $x \geq 3$  for which  $2x$  is never the sum of two primes, then for all

$p_1$  and  $p_2$  primes,  $3 \leq p_2 < p_1$ ,  $2x = p_1 + p_2$  OR  $2x = p_1 + p_2 + 2b_{p_1, p_2} = p_1 + p_2 + 2b$

then  $x = \frac{p_1 + p_2}{2} + b$ .

But for all  $p_1, p_2$  exists  $y$  for which  $y = \frac{p_1 - p_2}{2} + b$

Let

$$x_1 = p_1 + 2b, \quad x_2 = p_1 - 2b, \quad x_3 = p_2 + 2b, \quad x_4 = p_2 - 2b$$

We deduce that

$$x = \frac{p_1 + p_2}{2} + b = \frac{p_1 + x_2}{2} + b = \frac{x_1 + p_2}{2} = \frac{x_1 + x_2}{2} + b$$

$$= \frac{p_1 - x_2}{2} + \frac{x_1 + x_3}{2} + b = \frac{x_4 + x_2}{2} + 3b$$

$$y = \frac{p_1 - p_2}{2} + b = \frac{p_1 - x_2}{2} = \frac{x_1 - p_2}{2} = \frac{x_1 - x_2}{2} - b$$

$$= \frac{p_1 - x_3}{2} + 2b = \frac{x_1 - x_3}{2} + b = \frac{x_4 - x_3}{2} + 3b = \frac{x_4 - x_2}{2} + b$$

$$x_1 + x_2 = p_1 + p_2$$

$$x_1 - x_3 = p_1 - p_2$$

**Lemma 1**

The following formula

$$x = \frac{p_1 + p_2}{2} + b = \frac{p_1 + x_2}{2} + 2b = \frac{x_1 + p_2}{2} = \frac{x_1 + x_2}{2} + b$$

$$= \frac{p_1 + x_3}{2} = \frac{x_1 + x_3}{2} - b = \frac{x_4 + x_3}{2} + b = \frac{x_4 + x_2}{2} + 3b$$

$$y = \frac{p_1 - p_2}{2} + b = \frac{p_1 - x_2}{2} = \frac{x_1 - p_2}{2} = \frac{x_1 - x_2}{2} - b$$

$$= \frac{p_1 - x_3}{2} + 2b = \frac{x_1 - x_3}{2} + b = \frac{x_4 - x_3}{2} + 3b = \frac{x_4 - x_2}{2} + b$$

Imply that exist  $p_1$  and  $p_2$  prime numbers for which  $b = 0$

**Proof of lemma 1**

If  $x$  is prime  $2x = x + x$  is the sum of two primes, then  $p_1 - p_2 \neq 0$

We will suppose firstly that  $(x_1 - x_2)(x_1 + x_3) \neq 0$

Let

$$\frac{x_1 - x_2}{p_1 - p_2} = \frac{p_1 - p_2 + 4b}{p_1 - p_2} = 1 + \frac{4b}{p_1 - p_2}$$

$$\frac{p_1 - p_2}{x_1 - x_2} = \frac{x_1 - x_2 - 4b}{x_1 - x_2} = 1 - \frac{4b}{x_1 - x_2}$$

We pose  $k = \frac{2b}{p_1 - p_2}$ ,  $k' = -\frac{2b}{x_1 - x_2}$

$kk' = 0 \Rightarrow b = 0$ , we have supposed  $b \neq 0$

$\forall (x, y); \exists \varphi | x = \varphi y$

$x + y = (\varphi + 1)y = x_1 \neq 0$ ,  $x - y = (\varphi - 1)y = x_2 \neq 0$

$\forall (k, k'); \exists \alpha | k = \alpha k'$

$$\Rightarrow \frac{2b}{p_1 - p_2} = -\alpha \frac{2b}{x_1 - x_2}$$

$$\Rightarrow x_1 - x_2 = -\alpha \frac{p_1 - p_2}{x_1 - x_2} \Rightarrow x_1 - x_2 - p_1 + p_2 = 4b = -(\alpha + 1)(p_1 - p_2)$$

$$\Rightarrow b = -\frac{\alpha + 1}{4}(p_1 - p_2)$$

$$\Rightarrow x = \frac{p_1 + p_2}{2} + b = \frac{p_1 + p_2}{2} - \frac{\alpha + 1}{4}(p_1 - p_2) = \frac{(1 - \alpha)p_1 + (3 + \alpha)p_2}{4} = \frac{\varphi}{\varphi - 1} p_2$$

$$y = \frac{p_1 - p_2}{2} + b = \frac{p_1 - p_2}{2} - \frac{\alpha + 1}{4}(p_1 - p_2) = \frac{(1 - \alpha)(p_1 - p_2)}{4} = \frac{1}{\varphi - 1} p_2$$

Let

$$\frac{x_1 + x_3}{p_1 + p_2} = \frac{p_1 + p_2 + 4b}{p_1 + p_2} = 1 + \frac{4b}{p_1 + p_2}$$

$$\frac{p_1 + p_2}{x_1 + x_3} = \frac{x_1 + x_3 - 4b}{x_1 + x_3} = 1 - \frac{4b}{x_1 + x_3}$$

We pose  $m = \frac{2b}{p_1 + p_2}$ ,  $m' = -\frac{2b}{x_1 + x_3}$

$mm' = 0 \Rightarrow b = 0$ , we have supposed  $mm' \neq 0$

$\forall(m, m'); \exists \beta | m = \beta m'$

$$\Rightarrow \frac{2b}{p_1 + p_2} = -\beta \frac{2b}{x_1 + x_3}$$

$$\Rightarrow x_1 + x_3 = -\beta(p_1 + p_2) \Rightarrow x_1 + x_3 - p_1 - p_2 = 4b = -(\beta + 1)(p_1 + p_2)$$

$$\Rightarrow b = -\frac{\beta + 1}{4}(p_1 + p_2)$$

$$\Rightarrow x = \frac{p_1 + p_2}{2} + b = \frac{p_1 + p_2}{2} - \frac{\beta + 1}{4}(p_1 + p_2) = \frac{(1 - \beta)(p_1 + p_2)}{4} = \frac{\varphi}{\varphi - 1} p_2$$

$$y = \frac{p_1 - p_2}{2} + b = \frac{p_1 - p_2}{2} - \frac{\beta + 1}{4}(p_1 + p_2) = \frac{(1 - \beta)p_1 - (\beta + 3)p_2}{4} = \frac{1}{\varphi - 1} p_2$$

$$b = -\frac{\alpha + 1}{4}(p_1 - p_2) = -\frac{\beta + 1}{4}(p_1 + p_2) \Rightarrow (\beta - \alpha)p_1 = (-2 - \alpha - \beta)p_2$$

But :  $(2k+1)(2k'+1) = 1 \Rightarrow 2kk' + k + k' = 0 \Rightarrow (2k+1)k' = -k$

$$2k+1 = \frac{-k}{k'} = \frac{k+1}{k'+1} = \frac{a(k+1) + a'(2k+1)}{a(k'+1) + a'}$$

$$2k'+1 = \frac{-k'}{k} = \frac{k'+1}{k+1} = \frac{c(k'+1) + c'(2k'+1)}{c(k+1) + c'}, \quad \forall(a, a', c, c')$$

$$\frac{2k+1}{2k'+1} = \frac{((a+2a')k + a + a')(ck + c + c')}{((c+2c')k' + c + c')(ak' + a + a')}$$

$$= \frac{(ac + 2a'c)k^2 + (2ac + ac' + 3a'c + 2a'c')k + ac + ac' + a'c + a'c'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

$$\Rightarrow \frac{2k+1}{2k'+1} - 1 = (k - k') \frac{2}{2k'+1}$$

$$= \frac{ac(k^2 - k'^2) + 2a'ck^2 - 2ac'k'^2 + (2ac + ac' + a'c + 2a'c')(k - k') + 2a'c'k - 2ac'k'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

$\forall(a, a', c, c')$ , particularly  $(a, a', c, c') = (\delta k, a'c, \delta k, a'c)$

$$\Rightarrow (k - k') \frac{2}{2k'+1}$$

$$= \frac{ac(k+k')(k-k') + (2ac + ac' + a'c + 2a'c')(k - k') - 2\delta k^2 - 2\delta k^2 k'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

$$= (k - k') \frac{-2ac'k' + 2ac + ac' + a'c + 2a'c' - 2\delta k k'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

If  $k = k' \Rightarrow \alpha = 1 \Rightarrow \frac{(1 - \alpha)p_1 + (3 + \alpha)p_2}{4} = p_2$ , trivial solution : it is impossible.

$$k - k' \neq 0 \Rightarrow \frac{2}{2k'+1} = \frac{-2ac'k' + 2ac + ac' + a'c + 2a'c' - 2\delta k k'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

Also  $(2m+1)(2m'+1) = 1 \Rightarrow 2mm' + m + m' = 0 \Rightarrow (2m+1)m' = -m$

$$2m+1 = \frac{-m}{m'} = \frac{m+1}{m'+1} = \frac{a(m+1) + a'(2m+1)}{a(m'+1) + a'}$$

$$2m'+1 = \frac{-m'}{m} = \frac{m'+1}{m+1} = \frac{c(m'+1) + c'(2m'+1)}{c(m+1) + c'}, \quad \forall(a, a', c, c')$$

$$\frac{2m+1}{2m'+1} = \frac{((a+2a')m + a + a')(cm + c + c')}{((c+2c')m' + c + c')(am' + a + a')}$$

$$= \frac{(ac + 2a'c)m^2 + (2ac + ac' + 3a'c + 2a'c')m + ac + ac' + a'c + a'c'}{(ac + 2ac')m'^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$\Rightarrow \frac{2m+1}{2m'+1} - 1 = (m - m') \frac{2}{2m'+1}$$

$$= \frac{ac(m^2 - m'^2) + 2a'cm^2 - 2ac'm'^2 + (2ac + ac' + a'c + 2a'c')(m - m') + 2a'cm - 2ac'm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$\forall (a, a', c, c')$ , particularly  $(a, a', c, c') | ac' = \delta k^2 = \gamma m^2$ ,  $a'c = \delta k'^2 = \gamma' m'^2$

$$\Rightarrow (m - m') \frac{2}{2m'+1}$$

$$= \frac{ac(m+m')(m-m') + 2(\gamma' - \gamma)m^2m'^2 + (2ac + ac' + a'c + 2a'c')(m - m') + 2\gamma'mm'^2 - 2\gamma m^2m'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$= (m - m') \frac{-2acmm' + 2(\frac{\gamma' - \gamma}{m - m'})m^2m'^2 + 2ac + ac' + a'c + 2a'c' - 2(\frac{\gamma m - \gamma' m'}{m - m'})mm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$m = m' \Rightarrow \beta = 1 \Rightarrow x = \frac{(1 - \beta)(p_1 + p_2)}{4} = 0$ , it is impossible

But  $\frac{2}{2k'+1} = \frac{-2ackk' + 2ac + ac' + a'c + 2a'c' - 2\delta kk'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$

For  $(a, a', c, c') | ac' = \delta k^2 = \gamma m^2$ ,  $a'c = \delta k'^2 = \gamma' m'^2$

$$\Rightarrow \frac{2}{2m'+1} = \frac{-2acmm' + 2ac + ac' + a'c + 2a'c' - 2\gamma mm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$= \frac{-2acmm' + 2ac + ac' + a'c + 2a'c' - 2\gamma'mm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$\Rightarrow \frac{2}{2m'+1} = \frac{-2acmm' + 2(\frac{\gamma' - \gamma}{m - m'})m^2m'^2 + 2ac + ac' + a'c + 2a'c' - 2(\frac{\gamma m - \gamma' m'}{m - m'})mm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$= \frac{-2acmm' + 2ac + ac' + a'c + 2a'c' - 2\gamma mm'}{(ac + 2ac')m^2 + (2ac + 3ac' + a'c + 2a'c')m' + ac + ac' + a'c + a'c'}$$

$$= \frac{-2acmm' + 2(\frac{\gamma' - \gamma}{m - m'})m^2m'^2 + 2ac + ac' + a'c + 2a'c' - 2\gamma mm'}{(ac + 2ac')k'^2 + (2ac + 3ac' + a'c + 2a'c')k' + ac + ac' + a'c + a'c'}$$

$$\Rightarrow \gamma = \gamma' \Rightarrow ac' = \delta k^2 = \gamma m^2, a'c = \delta k'^2 = \gamma' m'^2 = \gamma m'^2$$

$$\Rightarrow \frac{\delta}{\gamma} = \frac{m^2}{k^2} = \frac{m'^2}{k'^2} \Rightarrow \frac{k^2}{k'^2} = (2k+1)^2 = \alpha^2 = \frac{m^2}{m'^2} = (2m+1)^2 = \beta^2$$

If  $\alpha = -\beta \Rightarrow (\beta - \alpha)p_1 = -(\beta + \alpha)p_2 = (-2 - \alpha - \beta)p_2 = -2p_2 \Rightarrow \alpha = -\beta = \frac{p_2}{p_1}$

$$b = -(\alpha + 1) \frac{p_1 - p_2}{4} - (\beta + 1) \frac{p_1 + p_2}{4} = -(\frac{\alpha + 1}{4} + \frac{\beta + 1}{4})(p_1 - p_2) = \frac{p_2^2 - p_1^2}{4p_1}$$

$\Rightarrow 4bp_1 = p_2^2 - p_1^2 \Rightarrow (4b + p_1)p_1 = p_2^2 \Rightarrow 4b + p_1 = \frac{p_2^2}{p_1}$  and it is impossible because  $p_1$  and  $p_2$  are primes

and  $\frac{p_2^2}{p_1}$  can not be an integer.

$$\Rightarrow \alpha - \beta = 0 \Rightarrow (\beta - \alpha)p_1 = (-2 - \alpha - \beta)p_2 = -2(1 + \alpha)p_2 = -2(1 + \beta)p_2 = 0 \Rightarrow \alpha = \beta = -1$$

$$\Rightarrow b = -\frac{(\alpha + 1)}{4}(p_1 - p_2) = -\frac{(\beta + 1)}{4}(p_1 + p_2) = 0$$

$$\Rightarrow x = \frac{p_1 + p_2}{2}, y = \frac{p_1 - p_2}{2}$$

Another proof : let  $u, u', v, v'$  verifying

$$\left(\frac{1}{up_1 - vp_2} - \frac{1}{u'p_1 + v'p_2}\right)p_1 = \left(\frac{1}{up_1 - vp_2} + \frac{1}{u'p_1 + v'p_2}\right)p_2$$

$$\Rightarrow (u' - u)p_1^2 + (v' + v)p_1p_2 = (u + u')p_1p_2 + (v' - v)p_2^2$$

Thus

$$(u'+u-v'-v)p_1p_2 + (u-u')p_1^2 = (v-v')p_2^2$$

We pose with a different of zero

$$u'+u-v'-v = a(\alpha - \beta)$$

$$\Rightarrow ((v-v') - a(2 + \alpha + \beta))p_2^2 = (u-u')p_1^2$$

Let

$$v-v'-a(2 + \alpha + \beta) = (2 + \alpha + \beta)^2$$

$$u-u' = (\alpha - \beta)^2$$

$$\Rightarrow u'p_1 + v'p_2 = (u - (\alpha - \beta)^2)p_1 + (v - a(2 + \alpha + \beta) - (2 + \alpha + \beta)^2)p_2$$

$$= up_1 + vp_2 - (\alpha - \beta)^2 p_1 - a(\alpha - \beta)p_1 - (2 + \alpha + \beta)(\alpha - \beta)p_1$$

Let  $up_1 = -vp_2$  hence

$$\left(\frac{1}{up_1 - vp_2} - \frac{1}{u'p_1 + v'p_2}\right)p_1 = \left(\frac{1}{up_1 - vp_2} + \frac{1}{u'p_1 + v'p_2}\right)p_2$$

$$\Rightarrow \left(\frac{1}{2u} + \frac{1}{(\alpha - \beta)^2 + a(\alpha - \beta) + (2 + \alpha + \beta)(\alpha - \beta)}\right)p_1$$

$$= \left(\frac{1}{2u} - \frac{1}{(\alpha - \beta)^2 + a(\alpha - \beta) + (2 + \alpha + \beta)(\alpha - \beta)}\right)p_2$$

$$\Rightarrow \frac{p_1 - p_2}{2u} = -p_1 \left(\frac{1}{(2 + \alpha + \beta)(2\alpha + a + 2)} + \frac{1}{(\alpha - \beta)(2\alpha + a + 2)}\right)$$

Let  $2u = \alpha - \beta$  hence

$$\frac{1}{\alpha - \beta} - \frac{1}{\alpha + \beta + 2} = -\frac{1}{(2 + \alpha + \beta)(2\alpha + a + 2)} - \frac{1}{(\alpha - \beta)(2\alpha + a + 2)}$$

$$\Rightarrow \frac{2\alpha + a + 3}{\alpha - \beta} = \frac{2\alpha + a + 1}{2 + \alpha + \beta}; \forall \alpha \Rightarrow \alpha = 0$$

It means  $b=0$  thus

$x + y = p_1, x - y = p_2$  and the primal  $y = r$  is the primal radius. As there is the condition  $p_2 < x < p_1$ ,

there is not an infinity of  $p_2, p_1$

$$\text{If } (x_1 - x_2)(x_1 + x_2) \Rightarrow (x_4 - x_3)(x_4 + x_3) = 0$$

$$\text{Let } \frac{x_4 + x_2}{p_1 + p_2} = \frac{p_1 + p_2 + 4b}{p_1 + p_2} = 1 + \frac{4b}{p_1 + p_2}$$

$$\frac{p_1 + p_2}{x_4 + x_2} = \frac{x_4 + x_2 + 4b}{x_4 + x_2} = 1 + \frac{4b}{x_4 + x_2}$$

$$k = -\frac{2b}{p_1 + p_2} = -\frac{2b}{x_4 + x_2}$$

$$\frac{x_4 - x_3}{p_1 - p_2} = \frac{p_1 - p_2 + 4b}{p_1 - p_2} = 1 + \frac{4b}{p_1 - p_2}$$

$$\frac{p_1 - p_2}{x_4 - x_3} = \frac{x_4 - x_3 + 4b}{x_4 - x_3} = 1 + \frac{4b}{x_4 - x_3}$$

$$m = -\frac{2b}{p_1 - p_2}, m' = \frac{2b}{x_4 - x_3}$$

With the same reasoning and calculus  $\Rightarrow b = 0$  But  $b$  can not be equal to zero in all cases, it means there is an impossibility related to the fact that the conjecture is undecidable and if it is so, it is true ! Because, we would find in the case it is undecidable and false with the computer the  $2x$  different of all sum of primes and it is contradictory !

Now, if we suppose that for all  $p_2, p_1$  primes, exists  $x \mid 2x \neq p_1 - p_2$

$x = \frac{p_1 - p_2}{2} + b, y = \frac{p_1 + p_2}{2} + b,$  with the same reasoning, the same calculus but replacing  $x$  by  $y$  and  $y$  by  $x$ , we prove that  $b = 0$ , which means that for all positive integer  $x$ , exists  $p_1, p_2$  for which  $x = \frac{p_1 - p_2}{2}$ , if we pose  $y = \frac{p_1 + p_2}{2}, x + y = p_1, y - x = p_2$

$y$  is the primal radius. As there is no condition on  $x, y, p_1, p_2$  there is an infinity of couples of primes  $(p_1, p_2)$  For  $x = 1, p_1$  and  $p_2$  are twin primes. Let us prove it. Let us suppose that exists an integer  $x \geq 0$  for which  $2x$  is never the difference of two primes, then for all  $p_1$  and  $p_2$  primes,  $p_2 < 2x + p_1 - p_2$ , or  $2x = p_1 - p_2 + 2b_{p_1, p_2} = p_1 - p_2 + 2b$ , then  $x = \frac{p_1 - p_2}{2} + b.$

$$y = \frac{p_1 + p_2}{2} + b$$

But for all  $p_1, p_2$  exists  $y$ , for which

Let

$$x_1 = p_1 + 2b, x_2 = p_2 - 2b, x_3 = p_2 + 2b, x_4 = p_1 - 2b$$

We deduce that

$$y = \frac{p_1 + p_2}{2} + b = \frac{p_1 + x_2}{2} + 2b = \frac{x_1 + p_2}{2} = \frac{x_1 + x_2}{2} + b$$

$$= \frac{p_1 + x_3}{2} = \frac{x_1 + x_3}{2} - b = \frac{x_4 + x_3}{2} + b = \frac{x_4 + x_2}{2} + 3b$$

$$x = \frac{p_1 - p_2}{2} + b = \frac{p_1 - x_2}{2} = \frac{x_1 - p_2}{2} = \frac{x_1 - x_2}{2} - b$$

$$= \frac{p_1 - x_3}{2} + 2b = \frac{x_1 - x_3}{2} + b = \frac{x_4 - x_3}{2} + 3b = \frac{x_4 - x_2}{2} + b$$

$$x_1 + x_2 = p_1 + p_2$$

$$x_1 - x_3 = p_1 - p_2$$

**Lemma 2**

The following formula

$$y = \frac{p_1 + p_2}{2} + b = \frac{p_1 + x_2}{2} + 2b = \frac{x_1 + p_2}{2} = \frac{x_1 + x_2}{2} + b$$

$$= \frac{p_1 + x_3}{2} = \frac{x_1 + x_3}{2} - b = \frac{x_4 + x_3}{2} + b = \frac{x_4 + x_2}{2} + 3b$$

$$x = \frac{p_1 - p_2}{2} + b = \frac{p_1 - x_2}{2} = \frac{x_1 - p_2}{2} = \frac{x_1 - x_2}{2} - b$$

$$= \frac{p_1 - x_3}{2} + 2b = \frac{x_1 - x_3}{2} + b = \frac{x_4 - x_3}{2} + 3b = \frac{x_4 - x_2}{2} + b$$

Imply that exist  $p_1$  and  $p_2$  prime numbers for which  $b = 0$

**Proof of lemma 2**

If  $x$  is prime  $0 = x - x$  is the sum of two primes, then  $p_1 - p_2 \neq 0$

We will suppose firstly that  $(x_1 - x_2)(x_1 + x_3) \neq 0$

Let

$$\frac{x_1 - x_2}{p_1 - p_2} = \frac{p_1 - p_2 + 4b}{p_1 - p_2} = 1 + \frac{4b}{p_1 - p_2}$$

$$\frac{p_1 - p_2}{x_1 - x_2} = \frac{x_1 - x_2 - 4b}{x_1 - x_2} = 1 - \frac{4b}{x_1 - x_2}$$

$$k = \frac{2b}{p_1 - p_2}, \quad k' = -\frac{2b}{x_1 - x_2}$$

We pose

$kk' = 0 \Rightarrow b = 0$ , we have supposed  $kk' \neq 0$

$\forall(x, y); \exists \varphi | y = \varphi x$

$x + y = (\varphi + 1)x = x_1 \neq 0, \quad y - x = (\varphi - 1)x = p_2 \neq 0$

$\forall(k, k'); \exists \alpha | k = \alpha k'$

$$\Rightarrow \frac{2b}{p_1 - p_2} = -\alpha \frac{2b}{x_1 - x_2}$$

$$\Rightarrow x_1 - x_2 = -\alpha(p_1 - p_2) \Rightarrow x_1 - x_2 - p_1 + p_2 = 4b = -(\alpha + 1)(p_1 - p_2)$$

$$\Rightarrow b = -\frac{\alpha + 1}{4}(p_1 - p_2)$$

$$\Rightarrow y = \frac{p_1 + p_2}{2} + b = \frac{p_1 + p_2}{2} - \frac{\alpha + 1}{4}(p_1 - p_2) = \frac{(1 - \alpha)p_1 + (3 + \alpha)p_2}{4} = \frac{\varphi}{\varphi - 1} p_2$$

$$x = \frac{p_1 - p_2}{2} + b = \frac{p_1 - p_2}{2} - \frac{\alpha + 1}{4}(p_1 - p_2) = \frac{(1 - \alpha)(p_1 - p_2)}{4} = \frac{1}{\varphi - 1} p_2$$

Let

$$\frac{x_1 + x_3}{p_1 + p_2} = \frac{p_1 + p_2 + 4b}{p_1 + p_2} = 1 + \frac{4b}{p_1 + p_2}$$

$$\frac{p_1 + p_2}{x_1 + x_3} = \frac{x_1 + x_3 - 4b}{x_1 + x_3} = 1 - \frac{4b}{x_1 + x_3}$$

$$m = \frac{2b}{p_1 + p_2}, \quad m' = -\frac{2b}{x_1 + x_3}$$

We pose

$mm' = 0 \Rightarrow b = 0$ , we have supposed  $mm' \neq 0$

$\forall(m, m'); \exists \beta | m = \beta m'$

$$\Rightarrow \frac{2b}{p_1 + p_2} = -\beta \frac{2b}{x_1 + x_3}$$

$$\Rightarrow x_1 + x_3 = -\beta(p_1 + p_2) \Rightarrow x_1 + x_3 - p_1 - p_2 = 4b = -(\beta + 1)(p_1 + p_2)$$

$$\Rightarrow b = -\frac{\beta + 1}{4}(p_1 + p_2)$$

$$\Rightarrow y = \frac{p_1 + p_2}{2} + b = \frac{p_1 + p_2}{2} - \frac{\beta + 1}{4}(p_1 + p_2) = \frac{(1 - \beta)(p_1 + p_2)}{4} = \frac{\varphi}{\varphi - 1} p_2$$

$$x = \frac{p_1 - p_2}{2} + b = \frac{p_1 - p_2}{2} - \frac{\beta + 1}{4}(p_1 + p_2) = \frac{(1 - \beta)p_1 - (\beta + 3)p_2}{4} = \frac{1}{\varphi - 1} p_2$$

$$b = -\frac{\alpha + 1}{4}(p_1 - p_2) = -\frac{\beta + 1}{4}(p_1 + p_2) \Rightarrow (\beta - \alpha)p_1 = (-2 - \alpha - \beta)p_2$$

But :  $(2k+1)(2k'+1)=1 \Rightarrow 2kk'+k+k'=0 \Rightarrow (2k+1)k'=-k$

$$2k+1 = \frac{-k}{k'} = \frac{k+1}{k'+1} = \frac{a(k+1)+a'(2k+1)}{a(k'+1)+a'}$$

$$2k'+1 = \frac{-k'}{k} = \frac{k'+1}{k+1} = \frac{c(k'+1)+c'(2k'+1)}{c(k+1)+c'}, \forall (a, a', c, c')$$

$$\frac{2k+1}{2k'+1} = \frac{((a+2a')k+a+a')(ck+c+c')}{((c+2c')k'+c+c')(ak'+a+a')}$$

$$= \frac{(ac+2a'c)k^2+(2ac+ac'+3a'c+2a'c')k+ac+ac'+a'c+a'c'}{(ac+2ac')k^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

$$\Rightarrow \frac{2k+1}{2k'+1} - 1 = (k-k') \frac{2}{2k'+1}$$

$$= \frac{ac(k^2-k'^2)+2a'ck^2-2ac'k'^2+(2ac+ac'+a'c+2a'c')(k-k')+2a'ck-2ac'k'}{(ac+2ac')k^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

$\forall (a, a', c, c')$ , particularly  $(a, a', c, c') | ac' = \delta k^2, a'c = \delta k'^2$

$$\Rightarrow (k-k') \frac{2}{2k'+1}$$

$$= \frac{ac(k+k')(k-k')+(2ac+ac'+a'c+2a'c')(k-k')+2\delta k k'^2-2\delta k'^2 k'}{(ac+2ac')k^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

$$= (k-k') \frac{-2ackk'+2ac+ac'+a'c+2a'c'-2\delta k k'}{(ac+2ac')k^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

$$k=k' \Rightarrow \alpha=1 \Rightarrow y = \frac{(1-\alpha)p_1+(3+\alpha)p_2}{4} = p_2 \text{ it is impossible.}$$

$$k-k' \neq 0 \Rightarrow \frac{2}{2k'+1} = \frac{-2ackk'+2ac+ac'+a'c+2a'c'-2\delta k k'}{(ac+2ac')k^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

Also  $(2m+1)(2m'+1)=1 \Rightarrow 2mm'+m+m'=0 \Rightarrow (2m+1)m'=-m$

$$2m+1 = \frac{-m}{m'} = \frac{m+1}{m'+1} = \frac{a(m+1)+a'(2m+1)}{a(m'+1)+a'}$$

$$2m'+1 = \frac{-m'}{m} = \frac{m'+1}{m+1} = \frac{c(m'+1)+c'(2m'+1)}{c(m+1)+c'}, \forall (a, a', c, c')$$

$$\frac{2m+1}{2m'+1} = \frac{((a+2a')m+a+a')(cm'+c+c')}{((c+2c')m'+c+c')(am'+a+a')}$$

$$= \frac{(ac+2a'c)m^2+(2ac+ac'+3a'c+2a'c')m+ac+ac'+a'c+a'c'}{(ac+2ac')m^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'}$$

$$\Rightarrow \frac{2m+1}{2m'+1} - 1 = (m-m') \frac{2}{2m'+1}$$

$$= \frac{ac(m^2-m'^2)+2a'cm^2-2ac'm'^2+(2ac+ac'+a'c+2a'c')(m-m')+2a'cm-2ac'm'}{(ac+2ac')m^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'}$$

$\forall (a, a', c, c')$ , particularly  $(a, a', c, c') | ac' = \delta k^2 = \gamma m^2, a'c = \delta k'^2 = \gamma' m'^2$

$$\Rightarrow (m-m') \frac{2}{2m'+1}$$

$$= \frac{ac(m+m')(m-m')+2(\gamma'-\gamma)m^2m'^2+(2ac+ac'+a'c+2a'c')(m-m')+2\gamma' mm'^2-2\gamma m^2 m'}{(ac+2ac')m^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'}$$

$$= (m-m') \frac{-2acmm'+2(\frac{\gamma'-\gamma}{m-m'})m^2m'^2+2ac+ac'+a'c+2a'c'-2(\frac{\gamma m-\gamma' m'}{m-m'})mm'}{(ac+2ac')m^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'}$$

$$m=m' \Rightarrow \beta=1 \Rightarrow y = \frac{(1-\beta)(p_1+p_2)}{4} = 0, \text{ it is impossible}$$



$$\text{But } \frac{2}{2k'+1} = \frac{-2ackk'+2ac+ac'+a'c+2a'c'-2\delta kk'}{(ac+2ac')k'^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'}$$

$$\text{For } (a, a', c, c') | ac' = \delta k^2 = \gamma m^2, \quad a'c = \delta k'^2 = \gamma' m'^2$$

$$\begin{aligned} \Rightarrow \frac{2}{2m'+1} &= \frac{-2acmm'+2ac+ac'+a'c+2a'c'-2\gamma mm'}{(ac+2ac')m'^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'} \\ &= \frac{-2acmm'+2ac+ac'+a'c+2a'c'-2\gamma' mm'}{(ac+2ac')m'^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'} \\ \Rightarrow \frac{2}{2m'+1} &= \frac{-2acmm'+2(\frac{\gamma'-\gamma}{m-m'})m^2m'^2+2ac+ac'+a'c+2a'c'-2(\frac{\gamma m-\gamma' m'}{m-m'})mm'}{(ac+2ac')m'^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'} \\ &= \frac{-2acmm'+2ac+ac'+a'c+2a'c'-2\gamma mm'}{(ac+2ac')m'^2+(2ac+3ac'+a'c+2a'c')m'+ac+ac'+a'c+a'c'} \\ &= \frac{-2acmm'+2(\frac{\gamma-\gamma'}{m-m'})m^2m'^2+2ac+ac'+a'c+2a'c'-2\gamma mm'}{(ac+2ac')k'^2+(2ac+3ac'+a'c+2a'c')k'+ac+ac'+a'c+a'c'} \end{aligned}$$

$$\Rightarrow \gamma = \gamma' \Rightarrow ac' = \delta k^2 = \gamma m^2, \quad a'c = \delta k'^2 = \gamma' m'^2 = \gamma m'^2$$

$$\Rightarrow \frac{\delta}{\gamma} = \frac{m^2}{k^2} = \frac{m'^2}{k'^2} \Rightarrow \frac{k^2}{k'^2} = (2k+1)^2 = \alpha^2 = \frac{m^2}{m'^2} = (2m+1)^2 = \beta^2$$

$$\text{If } \alpha = -\beta \Rightarrow (\beta - \alpha)p_1 = -2\alpha p_1 = 2\beta p_1 = (-2 - \alpha - \beta)p_2 = -2p_2 \Rightarrow \alpha = -\beta = \frac{p_2}{p_1}$$

$$b = -(\alpha+1)\frac{p_1-p_2}{4} = -(\beta+1)\frac{p_1+p_2}{4} = -\frac{(p_1+p_2)(p_1-p_2)}{4p_1} = \frac{p_2^2-p_1^2}{4p_1}$$

$$\Rightarrow 4bp_1 = p_2^2 - p_1^2 \Rightarrow (4b+p_1)p_1 = p_2^2 \Rightarrow 4b+p_1 = \frac{p_2^2}{p_1} \text{ and it is not possible because } p_1 \text{ and } p_2 \text{ are primes}$$

$$\text{and } 4b+p_1 = \frac{p_2^2}{p_1} \text{ can not be an integer.}$$

$$\Rightarrow \alpha - \beta = 0 \Rightarrow (\beta - \alpha)p_1 = (-2 - \alpha - \beta)p_2 = -2p_2 + \beta p_2 = 0 \Rightarrow \alpha = \beta = -1$$

$$\Rightarrow b = -\frac{(\alpha+1)}{4}(p_1-p_2) = -\frac{(\beta+1)}{4}(p_1+p_2) = 0$$

$$\Rightarrow y = \frac{p_1+p_2}{2}, \quad x = \frac{p_1-p_2}{2}$$

Another proof: let  $u, u', v, v'$  be solving

$$\left(\frac{1}{up_1-vp_2} - \frac{1}{u'p_1+v'p_2}\right)p_1 = \left(\frac{1}{up_1-vp_2} + \frac{1}{u'p_1+v'p_2}\right)p_2$$

$$\Rightarrow (u'-u)p_1^2 + (v'-v)p_1p_2 = (u+u')p_1p_2 + (v'-v)p_2^2$$

Thus

$$(u'+u-v'-v)p_1p_2 + (u-u')p_1^2 = (v-v')p_2^2$$

We pose  $a$  a different of zero

$$u'+u-v'-v = a(\alpha - \beta)$$

$$\Rightarrow ((v-v')-a(2+\alpha+\beta))p_2^2 = (u-u')p_1^2$$

Let

$$v-v'-a(2+\alpha+\beta) = (2+\alpha+\beta)^2$$

$$u-u' = (\alpha - \beta)^2$$

$$\Rightarrow u'p_1 + v'p_2 = (u - (\alpha - \beta)^2)p_1 + (v - a(2 + \alpha + \beta) - (2 + \alpha + \beta)^2)p_2$$

$$= up_1 + vp_2 - (\alpha - \beta)^2 p_1 - a(\alpha - \beta)p_1 - (2 + \alpha + \beta)(\alpha - \beta)p_1$$

Let  $up_1 = -vp_2$  hence

$$\begin{aligned} \left(\frac{1}{up_1 - vp_2} - \frac{1}{u'p_1 + v'p_2}\right)p_1 &= \left(\frac{1}{up_1 - vp_2} + \frac{1}{u'p_1 + v'p_2}\right)p_2 \\ \Rightarrow \left(\frac{1}{2u} + \frac{1}{(\alpha - \beta)^2 + a(\alpha - \beta) + (2 + \alpha + \beta)(\alpha - \beta)}\right)p_1 \\ &= \left(\frac{1}{2u} - \frac{1}{(\alpha - \beta)^2 + a(\alpha - \beta) + (2 + \alpha + \beta)(\alpha - \beta)}\right)p_2 \\ \Rightarrow \frac{p_1 - p_2}{2u} &= -p_1 \left(\frac{1}{(2 + \alpha + \beta)(2\alpha + a + 2)} + \frac{1}{(\alpha - \beta)(2\alpha + a + 2)}\right) \end{aligned}$$

Let  $2u = \alpha - \beta$  hence

$$\begin{aligned} \frac{1}{\alpha - \beta} - \frac{1}{\alpha + \beta + 2} &= -\frac{1}{(2 + \alpha + \beta)(2\alpha + a + 2)} - \frac{1}{(\alpha - \beta)(2\alpha + a + 2)} \\ \Rightarrow \frac{2\alpha + a + 3}{\alpha - \beta} &= \frac{2\alpha + a + 1}{2 + \alpha + \beta}; \forall a \Rightarrow \alpha = \beta = -1 \end{aligned}$$

It means  $b=0$  thus

$x + y = p_1, y - x = p_2$  are primes  $y = r$  is the primal radius. As there is no condition, there is an infinity of  $p_1, p_2$ .

If  $(x_1 - x_2)(x_1 + x_2) = 0 \Rightarrow (x_4 + x_2)(x_4 - x_3) \neq 0$

$$\text{Let } \frac{x_4 + x_2}{p_1 + p_2} = \frac{p_1 + p_2 - 4b}{p_1 + p_2} = 1 - \frac{4b}{p_1 + p_2}$$

$$\frac{p_1 + p_2}{x_4 + x_2} = \frac{x_4 + x_2 + 4b}{x_4 + x_2} = 1 + \frac{4b}{x_4 + x_2}$$

$$k = -\frac{2b}{p_1 + p_2}, k' = \frac{2b}{x_4 + x_2}$$

$$\frac{x_4 - x_3}{p_1 - p_2} = \frac{p_1 - p_2 - 4b}{p_1 - p_2} = 1 - \frac{4b}{p_1 - p_2}$$

$$\frac{p_1 - p_2}{x_4 - x_3} = \frac{x_4 - x_3 + 4b}{x_4 - x_3} = 1 + \frac{4b}{x_4 - x_3}$$

$$m = -\frac{2b}{p_1 - p_2}, m' = \frac{2b}{x_4 - x_3}$$

With the same calculus and reasoning, it implies that  $b = 0$ . But  $b$  can not be equal to zero in all cases, it means there is an impossibility related to the fact that the conjecture is undecidable and if it is so, it is true! Because, we could find in the case it is undecidable and false with the computer the  $2x$  difference of all sum of primes and it is contradictory!

For  $2x = p_1 - p_2$  is a difference of an infinity of couples of primes. There is an infinity of consecutive primes. And for all  $x > 2104$  exists  $p_1 = p_2 + 2, p_2$  primes for which  $2x = p_1 + p_2$

### Conclusion

The notion of the primal radius as defined in this study allows to confirm that for all integer  $x \geq 3$  exists a number  $r > 0$  for which  $x + r$  and  $x - r$  are primes and that for all integer  $x \geq 0$  exists a number  $r > 0$  for which  $x + r$  and  $r - x$  are primes and that exists an infinity of such primes.  $r$  is called the primal radius. The corollary is the proof of the Goldbach conjecture and de Polignac conjecture which stipulate, the first that an even number is always the sum of two prime numbers, the second that an even number is always the difference between two primes and that there is an infinity of such couples of primes. Another corollary is the proof of the twin primes conjecture which stipulates that there is an infinity of consecutive primes.

---

**References**

- [1] J. R. Chen, 2002, On the representation of a larger even integer as the sum of a prime and the product of at most two primes. *Sci. Sinica* 16, 157-176.
- [2] D. R. Heath-Brown, J. C. Puchta, 2002, Integers represented as a sum of primes and powers of two. *The Asian Journal of Mathematics*, 6, no. 3, pages 535-565.
- [3] H.L. Montgomery, Vaughan, R. C., 1975, The exceptional set in Goldbach's problem. Collection of articles in memory of Jurii Vladimirovich Linnik. *Acta Arith.* 27, 353–370.
- [4] J. Richstein, 2001, Verifying the goldbach conjecture up to  $4 \cdot 10^{14}$ , *Math. Comp.*, 70:236 ,1745--1749.
- [5] L. E. Dickson, 2005, *History of The Theory of Numbers*, Vol1, New York Dover.

**RETRACTED**