OBSERVATIONS ON THE TERNARY QUADRATIC EQUATION

$$X^2 = 24\alpha^2 + Y^2$$

M.A.Gopalan¹, V.Sangeetha², Manju Somanath³

¹Professor, Department of Mathematics, Srimathi Indira Gandhi College, Trichy-2, India.

^{2,3}Assistant Professor, Department of Mathematics, National College, Trichy-1, India.

Keywords. Ternary quadratic, polygonal numbers, integral solutions.

Abstract. The ternary quadratic equation $X^2 = 24\alpha^2 + Y^2$ is considered. Employing its non-zero integral solutions, relations among a few special polygonal numbers are determined.

1. INTRODUCTION

In [1,8,10], different patterns of m-gonal numbers are presented. In [2], explicit formulas for the ranks of triangular numbers which are simultaneously equal to pentagonal, decagonal and dodecagonal in turn are presented. In [3,5,6], the relations among the pairs of special m-gonal numbers are generated through the solutions of the ternary quadratic equation are determined. In [4,7,9], the binary quadratic equations are considered and relations between special polygonal numbers are obtained through its solutions.

In this communication, we consider the ternary quadratic equation given by $X^2 = 24\alpha^2 + Y^2$, where α is a non-zero integer and obtain the relations among the pairs of special m-gonal numbers generated through its solutions.

2. NOTATION

- $t_{3,N}$ · Triangular number of rank N
- $t_{5,P}$ · Pentagonal number of rank P
- $t_{6,Q}$ · Hexagonal number of rank Q
- $t_{7,H}$ · Heptagonal number of rank H
- $cp_{8,M}$ · Centered octagonal number of rank M
- $g_4(G)$ · 4-gram number of rank G

3. METHOD OF ANALYSIS

Consider the Diophantine equation

$$X^2 = 24\alpha^2 + Y^2 \quad (1)$$

where α is a non-zero integer. The smallest non-zero integer solution of the Pellian equation $X^2 = 24\alpha^2 + 1$ is given by $\alpha_0 = 1$, $X_0 = 5$. Thus the general form of non-trivial integral solutions (α_n, X_n) are given by

$$\begin{aligned} \alpha_n &= \frac{Y}{4\sqrt{6}} \left[(5 + 2\sqrt{6})^{n+1} - (5 - 2\sqrt{6})^{n+1} \right], \quad n = 0, 1, 2, \dots \\ X_n &= \frac{Y}{2} \left[(5 + 2\sqrt{6})^{n+1} + (5 - 2\sqrt{6})^{n+1} \right] \end{aligned} \quad (2)$$

Case 3.1

Let N and Q be the ranks of the n^{th} Triangular and Hexagonal numbers respectively.

The choice

$$X = 2N + 1; Y = 4Q - 1 \quad (3)$$

in (1) leads to the relation “ 2 times triangular number – 2 times hexagonal number = a nasty number” From (2) and (3), the values of ranks of Triangular numbers are found to be

$$N_n = \frac{4Q-1}{4} \left[(5 + 2\sqrt{6})^{n+1} + (5 - 2\sqrt{6})^{n+1} \right] - \frac{1}{2}, \quad n = 0, 1, 2, \dots \text{ and } Q \in \mathbb{Z}^+ - \{0\}$$

Numerical examples are illustrated in the table 3.1

Table 3.1

Q	$t_{6,Q}$	N_n $n = 0, 1, 2, \dots$	t_{3,N_n} $n = 0, 1, 2, \dots$
1	1	7,73,727,....	28,2701,264628,....
2	6	17,171,1697,...	153,14706,1440753,....
3	15	27,269,2667,...	378,36315,3557778,...

Case 3.2

Let N and P be the ranks of the n^{th} Triangular and Pentagonal numbers respectively.

The choice

$$X=2N+1; Y=6P-1 \quad (4)$$

in (1) leads to the relation “ 2 times triangular number – 6 times pentagonal number = a nasty number” From (2) and (4), the values of ranks of Triangular numbers are found to be

$$N_n = \frac{6P-1}{4} \left[(5 + 2\sqrt{6})^{n+1} + (5 - 2\sqrt{6})^{n+1} \right] - \frac{1}{2}, \quad n = 0, 1, 2, \dots \text{ and } P \in \mathbb{Z}^+ - \{0\}$$

Numerical examples are illustrated in the table 3.2

Table 3.2

P	$t_{5,P}$	N_n $n = 0, 1, 2, \dots$	t_{3,N_n} $n = 0, 1, 2, \dots$
1	1	12,122,1212,....	78,7503,735078,....
2	5	27,269,2667,...	378,36315,3557778,...
3	12	42,416,4122,...	903,86736,8497503,....

Case 3.3

Let N and H be the ranks of the n^{th} Triangular and Heptagonal numbers respectively.

The choice

$$X=6N+3; Y=10H-3 \quad (5)$$

in (1) leads to the relation “ 18 times triangular number – 10 times heptagonal number = a nasty number” From (2) and (5), the values of ranks of Triangular numbers are found to be

$$N_n = \frac{10H-3}{12} \left[(5 + 2\sqrt{6})^{n+1} + (5 - 2\sqrt{6})^{n+1} \right] - \frac{1}{2}, \quad n = 0, 1, 2, \dots \text{ and } H \equiv 0 \pmod{3}$$

Numerical examples are illustrated in the table 3.3

Table 3.3

H	$t_{7,H}$	N_n $n = 0, 1, 2, \dots$	t_{3,N_n} $n = 0, 1, 2, \dots$
3	18	22,220,2182,...	253,24310,2381653,....
6	81	47,465,4607,...	1128,108345,10614528,....
9	189	72,710,7032,...	2628,252405,24728028,...

Case 3.4

Let M and G be the ranks of the n^{th} Centered octagonal and 4-gram numbers respectively.

The choice

$$X=8M+3; Y=8G-4 \quad (6)$$

in (1) leads to the relation “4 times centered octagonal number – 4 times 4-gram number = a nasty number” From (2) and (6), the values of ranks of centered octagonal numbers are found to be

$$N_n = \frac{8G-4}{16} \left[(5+2\sqrt{6})^{n+1} + (5-2\sqrt{6})^{n+1} \right] - \frac{1}{2}, \quad n = 0, 1, 2, \dots \text{ and } G \in \mathbb{Z}^+ - \{0\}$$

Numerical examples are illustrated in the table 3.4

Table 3.4

G	$g_4(G)$	M_n $n = 0, 1, 2, \dots$	cp_{8, M_n} $n = 0, 1, 2, \dots$
1	1	2, 24, 242,	25, 2401, 235225,
2	9	7, 73, 727, ...	225, 21609, 2117025, ...
3	25	12, 122, 1212, ...	625, 60025, 5880625,

CONCLUSION

To conclude, one may determine the relations between the special polygonal numbers employing the integer solutions of various ternary quadratic equations.

REFERENCES

- [1] Dickson. L E, History of theory of numbers, Chelsea Publishing Company, New York (2) (1971).
- [2] Gopalan M A and Devibala S, Equality of triangular numbers with special m-gonal numbers, Bulletin of Allahabad Mathematical Society, 1(2006), 25-29.
- [3] Gopalan M A., Manju Somanath and Vanitha N, Observations on $X^2=8\alpha^2+Y^2$, Advances in Theoretical and Mathematical Society, 3 (2006), 245-248.
- [4] Gopalan M A and Srividhya G, Relations among m-gonal numbers through the equation $y^2=2x^2-I$, Antarctica J.Math, 7(3) (2010), 363-369.
- [5] Gopalan. M A and Srividhya G, Observations on $X^2=48\alpha^2+Y^2$, Archimedes J.Math., 2(1) (2012), 1-5.
- [6] Gopalan. M A and Srividhya G, Observations on $Y^2=2X^2+Z^2$, Archimedes J.Math., 2(1) (2012), 1-5.
- [6] Gopalan. M A and Srividhya G, Observations on , Archimedes J.Math., 2(1) (2012), 7-15
- [7] Gopalan. M A and Geetha V, Observations on some Pellian equations, Cayley J.Math., 2(2) (2013), 109-118.
- [8] Kapur J N, Ramanujan's Miracles, Mathematical Sciences Trust Society (1997).
- [9] Manju Somanath, Gopalan M A and Sangeetha G, Relations among special figurate numbers through the equation $y^2=10x^2+I$, Impact J.Sci.Tech, 5(1) (2011), 57-60.
- [10] Shailesh Shirali, Mathematical Marvels, A primer on number sequences, University Press (2001).