Lanczos approach to Noether's theorem

P. Lam-Estrada¹, J. López-Bonilla², R. López-Vázquez²

¹ Dept. of Maths., National Polytechnic Institute (IPN), Edif. 9, Col. Lindavista CP 07738, México

² ESIME-Zacatenco, IPN, Edif. 5, 1er. Piso, Col. Lindavista CP 07738, Mexico city;

jlopezb@ipn.mx

Keywords: Noether's theorem; Invariance of the action; Variational Lanczos technique.

Abstract. If the action $A = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt$ is invariant under the infinitesimal transformation $\tilde{t} = t + \varepsilon \tau(q, t)$, $\tilde{q}_r = q_r + \varepsilon \zeta_r(q, t)$, $r = 1, \dots, n$, with $\varepsilon = \text{constant} \ll 1$, then the Noether's theorem permits to construct the corresponding conserved quantity. The Lanczos method accepts that $\varepsilon = q_{n+1}$ is a new degree of freedom, thus the Euler-Lagrange equation for this new variable gives the Noether's constant of motion.

If in the functional (the concept of action was proposed by Leibnitz [1]) $A = \int_{t_1}^{t_2} L(q, \dot{q}, t) dt$ we apply the infinitesimal transformation ($\varepsilon = \text{constant} \ll 1$):

$$\tilde{t} = t + \varepsilon \tau(q, t), \quad \tilde{q}_r = q_r + \varepsilon \zeta_r(q, t), \quad r = 1, \dots, n \quad (1)$$

that is

$$\tilde{A} = \int_{\tilde{t}_1}^{\tilde{t}_2} L(\tilde{q}, \frac{d\tilde{q}}{d\tilde{t}}, \tilde{t}) d\tilde{t} \quad (2)$$

then we say that the action is invariant if:

$$\tilde{A} = A + \varepsilon \int_{t_1}^{t_2} \frac{d}{dt} F(q, t) dt, \quad (3)$$

thus the Euler-Lagrange equations (Lagrangian expressions [2, 3]) corresponding to the variational principle $\delta A = 0$:

$$E_r \equiv \frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_r} \right) - \frac{\partial L}{\partial q_r} = 0, \quad r = 1, \dots, n \quad (4)$$

remain intact. Noether [2] studied the case $F = 0$, and she suggested [3, 4] to Bessel-Hagen [5] the analysis of (4) with $F \neq 0$.

Therefore, we have a symmetry up to divergence and Noether [2, 5-8] proved the existence of the Rund-Trautman identity [6, 7, 9, 10]:

$$\frac{\partial L}{\partial q_r} \zeta_r + \frac{\partial L}{\partial \dot{q}_r} \dot{\zeta}_r + \frac{\partial L}{\partial t} \tau - \left(\frac{\partial L}{\partial \dot{q}_r} \dot{q}_r - L \right) \dot{\tau} - \frac{dF}{dt} = 0, \quad (5)$$

which can be written in the form:

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{q}_r} \zeta_r - H\tau - F \right) = (\zeta_r - \dot{q}_r \tau) E_r, \quad H = \frac{\partial L}{\partial \dot{q}_c} \dot{q}_c - L. \quad (6)$$

In (5) and (6) we employ the convention of Dedekind [11, 12]-Einstein because we sum over repeated indices.

If in (6) we use the Euler-Lagrange equations (4) we deduce the constant of motion associated to (1):

$$\frac{\partial L}{\partial \dot{q}_r} \zeta_r - H\tau - F = \text{Constant}, \quad (7)$$

thus we have a connection between symmetries and conservation laws [3, 6, 7, 9, 13-15].

In the next Section we show the Lanczos technique [16-18] to obtain the Noether's conserved quantity (7).

Lanczos method to construct conservation laws

Lanczos [16, 17] applies the infinitesimal transformation (1) (with $\varepsilon = \text{constant}$) to the action (2) and uses expansion of Taylor up to first order in ε , thus:

$$\tilde{A} = A + \varepsilon \int_{t_1}^{t_2} \left(\frac{\partial L}{\partial q_r} \zeta_r + \frac{\partial L}{\partial \dot{q}_r} \dot{\zeta}_r + \frac{\partial L}{\partial t} \tau - H\tau \right) dt,$$

then this integrand is equal to $\frac{dF}{dt}$ in harmony with the Rund-Trautman identity (5).

Now Lanczos proposes to employ (1) into (2) but considering that ε is a function, therefore up to 1th order in ε :

$$\tilde{A} = \int_{t_1}^{t_2} \left[L + \varepsilon \frac{dF}{dt} + \dot{\varepsilon} \left(\frac{\partial L}{\partial \dot{q}_r} \zeta_r - H\tau \right) \right] dt = \int_{t_1}^{t_2} \bar{L} dt, \quad (8)$$

$$\frac{d}{dt} \left(\frac{\partial \bar{L}}{\partial \dot{\varepsilon}} \right) - \frac{\partial \bar{L}}{\partial \varepsilon} = 0. \quad (9)$$

It is clear that:

$$\frac{\partial \bar{L}}{\partial \dot{\varepsilon}} = \frac{\partial L}{\partial \dot{q}_r} \zeta_r - H\tau, \quad \frac{\partial \bar{L}}{\partial \varepsilon} = \frac{dF}{dt},$$

therefore (9) implies (7). In other words, if the parameter of the symmetry is considered as an additional degree of freedom of the variational principle, then its Euler-Lagrange equation gives the Noether's constant of motion.

The work [18] has applications of this Lanczos technique to some Lagrangians employed in [8, 19-21].

References

- [1]. G. Leibnitz, *Dynamica de potentia et leqibus nature corporae* (1669) (published in 1890)
- [2]. E. Noether, *Invariante variationsprobleme*, *Nachr. Ges. Wiss. Göttingen* 2 (1918) 235-257
- [3]. Y. Kosmann-Schwarzbach, *The Noether theorems*, Springer, New York (2011)
- [4]. P. Havas, *The connection between conservation laws and invariance groups: folklore, fiction and fact*, *Acta Physica Austriaca* 38 (1973) 145-167
- [5]. E. Bessel-Hagen, *Über die erhaltungssätze der elektrodynamik*, *Mathematische Annalen* 84 (1921) 258-276
- [6]. D. E. Neuenschwander, *Symmetries, conservation laws, and Noether's theorem*, *Radiations, fall* (1998) 12-15
- [7]. D. E. Neuenschwander, *Emmy Noether's wonderful theorem*, The Johns Hopkins University Press, Baltimore (2011)
- [8]. M. Havelková, *Symmetries of a dynamical system represented by singular Lagrangians*, *Comm. in Maths* 20, No. 1 (2012) 23-32
- [9]. A. Trautman, *Noether's equations and conservation laws*, *Commun. Math. Phys.* 6 (1967) 248-261
- [10]. H. Rund, *A direct approach to Noether's theorem in the calculus of variations*, *Utilitas Math.* 2 (1972) 205-214
- [11]. M. A. Sinaceur, *Dedekind et le programme de Riemann*, *Rev. Hist. Sci.* 43 (1990) 221-294
- [12]. D. Laugwitz, *Bernhard Riemann 1826-1866. Turning points in the conception of mathematics*, Birkhäuser, Boston MA (2008)

-
- [13]. N. Byers, Emmy Noether's discovery of the deep connection between symmetries and conservation laws, Proc. Symp. Heritage E. Noether, Bar Ilan University, Tel Aviv, Israel, 2-3 Dec. (1996)
- [14]. K. A. Brading, H. R. Brown, Symmetries and Noether's theorems, in "Symmetries in Physics, Philosophical Reflections", Eds. Katherine A. Brading, Elena Castellani, Cambridge University Press (2003) 89-109
- [15]. L. M. Lederman, Ch. T. Hill, Symmetry and the beautiful Universe, Prometheus Books, Amherst, New York (2004) Chaps. 3 and 5
- [16]. C. Lanczos, The variational principles of mechanics, University of Toronto Press (1970) Chap. 11
- [17]. C. Lanczos, Emmy Noether and the calculus of variations, Bull. Inst. Math. and Appl. 9, No. 8 (1973) 253-258
- [18]. P. Lam-Estrada, J. López-Bonilla, R. López-Vázquez, G. Ovando, Lagrangians: Symmetries, gauge identities and first integrals, The SciTech, J. of Sci. & Tech. 3, No. 1 (2014) 54-66
- [19]. M. Henneaux, C. Teitelboim, J. Zanelli, Gauge invariance and degree of freedom count, Nucl. Phys. B332, No. 1 (1990) 169-188
- [20]. H. J. Rothe, K. D. Rothe, Classical and quantum dynamics of constrained Hamiltonian systems, World Scientific Lecture Notes in Physics 81, Singapore (2010)
- [21]. G. F. Torres del Castillo, Point symmetries of the Euler-Lagrange equations, Rev. Mex. Fis. 60 (2014) 129-135