PARTICLE KNOTS IN TORIC MODULAR SPACE

J.A.de Wet

Box 514,Plettenberg Bay,6600,South Africa

Keywords: Compactification, Coxeter Graphs, GF(4),Torus Knots, QCD, Triality.

Abstract

The goal of this contribution is to relate quarks to knots or loops in a 6-space CP^3 that then ollapses into a torus in real 3-space P^3 instantaneously after the Big Bang, and massive inflation, when 3 quarks unite to form nucleons.

Introduction

Kedia et. al. in recent paper [10] investigate knotted structures in hydrodynamic fields such as current-guiding magnetic field lines in a plasma, or vortex lines of classical or quantum fluids which arise naturally as excitations that carry helicity that is a measure of the knottedness of the field. In particular their Fig.2g is a trefoil which is our Fig.3 without the quadrupole (that will be seen in Section 2 to collapse into a point) and the color-coding.

Fig. 1

Fig. 2

Fig. 3

The torus shown in Fig.2 is due to Marcelis [13] whose calculations in a projective space with 24 vertices appear to be unpublished, but are supported to some extent by Westy [18] (from the same school) who provides a color-coded complex map of the Riemann surface z that incorporates the phases of $\omega=120$ degrees.

Murasagi [14] Ch.7 shows that Fig.2 is a trefoil (3,2) on a torus when we choose 3 points on the ends of a cylinder that can be joined to form the torus. This brings us to the goal of this contribution which is to relate the elementary particles to knots or loops in a 6-space. Here we will be guided by the work of Coxeter [5,6] who specifically labels the vertices of the torus appearing in Fig.1 by $0, \pm 1, \omega, \omega^2$ where $\omega = \exp(2i\pi/3)$ so that a knot crosses the longitude of a torus at $\omega=120$ degrees. Essentially this is a Galois Field GF(4) with permutations of ω raised to the powers 0,1,2,3 that will be considered in more detail in the next Section where we will show how 3 quarks in 6-space unite to become a nucleon in the projective space CP^3 which collapses to P^3 immediately after the Big Bang. Section 3 will employ the color-coding of Fig.1 as a model for Quantum Chromodynamics or QCD. Finally according to Rovelli [15] knots or loops in the 6-space described by E_6 employed by Coxeter may also describe Loop Quantum Gravity although details are beyond the scope of this contribution. Also Arvin [2] has considered knots on a torus as a model for elementary particles but excluding quarks. Again, Sundance O Billson Thompson, Smolin et.al. [17] also use knots as a model for Quantum Gravity and the Standard Model but utilize trinions instead of trefoils.

Coxeter Algebra

Fig.1 is a torus taken from Coxeter [5] which is an alternative to the graph of $su(3)_{\text{color}} \times su(3)_{\text{spin}} \times su(3)_{\text{isospin}}$ which is a triality sub-algebra of E_6 . This was published later [6] as Fig.12.3B. Both graphs are orbifolds with 27 vertices that according to Slansky [16] may be labeled by particles in the Standard Model or SM. However the actual labeling of the tritangent planes on a cubic surface (discussed by Hunt [10] Ch. 4) is new, but in line with Coxeter's labels. For example the up-quark u in Fig.1 is labeled by (012) indicated by $0, \omega, \omega^2$. Thus (023) on the same tritangent is simply a rotation through $\omega=120$ degrees and so on. In this way we find an equilateral triangle labeled by the 3 quarks uud comprising a proton and another ddu for the neutron beginning with (120). There are 2 more tritangents (not labeled) for the anti-particles which complete the outer ring of Fig.1. But quarks also belong to a GF(4) ring and thus the trefoil on a torus which is precisely the model adopted by Green, Schwarz and Witten [8] Section 9.5.2.

The quarks at the vertices of Fig.1 are trefoils illustrated by Fig.2, but the torus in Fig.1 only becomes a trefoil after the collapse of the inner ring just after the Big Bang when quarks in the 6-space CP^3 unite to build nucleons in the projective space P^3 .

This is supported by Barina and Jureto [5] where only the 12 outer vertices and the center of Fig.1 are in P^3 . Specifically these authors find 15 synthemes, where a syntheme has 6 'fix-lines' that are the edges of an invariant tetrahedron such as $u u d 0$ representing a proton. However because there are only 3 vertices on the face of a syntheme the outer ring of Fig.1 carries the 4 stable particles proton, neutron and their anti-particles. Also since the tritangents are invariant under rotations there are actually $3 \times 4 = 12$ possible synthemes on the outer ring. Specifically each syntheme consists of 3 commuting operators. Thus 3 synthemes can be chosen for spin rotations about the 3 axes of 3-space. Thus introducing triality which is a characteristic of Toric-Calabi-Yau moduli space that carry the Hessian Polyhedra in E_6 as discussed by Lie-Yang [12] and analysed by Coxeter [5].

Specifically each rotation in 3-space is also accompanied by a corresponding rotation in a parity 4-space when we permute 1,2,3. Charge conjugation belongs to a second set of 3 synthemes with the same rotations in a 3-space but a parity 5-space in another charge space [7].

In this way the 12 unstable particles $sss, \overline{sss}, \mu^\pm, \nu_\mu; \tau^\pm, \nu_\tau$ do not appear in the compactification of CP^3 to P^3 . This may be visualised as a collapse of the inner vertices to a point which carries the remaining 3 synthemes $e^\pm, \nu_e$ labeled by $\{011, 022, 033\}, \{110, 220, 330\}$ and $\{101, 202, 303\}$ for the muon. In this process the masses m_τ, m_ν of the τ, μ reappear as stable deuterium 3

according to the relationship

$$m_\tau + m_\mu = m_p + m_n + m_e \quad (1)$$

There is no heavy-ion decay and the same relation holds for the anti-particles. This equation is accurate if we assume that $m_\tau = 1777$ MeV and $m_\mu = 101.4$ MeV instead of the Fermi decomposition of muon decay in the weak interaction yielding 106 MeV. However in a recent publication Benjamin Brau et.al.[4] find a value of approximately 100 MeV for the mass of cosmic-ray muons so there is as yet some experimental uncertainty.

Quantum Chromodynamics,QCD

Returning again to Fig.1,when the inner vertices are contracted to a point at the origin the red, green and blue lines could serve as gluons on a new torus where a red upper path passes through the center before emerging at the circumference and giving way to a green gluon that in turn passes under the torus and then over to connect with a down quark and so on.The 3 color complex dimensions vanish when $CP^3 \rightarrow P^3$ but a torus knot remains in 3-space.

However Marcelis [13] calculates the dual set of 3 paths for the anti-gluons overline(r,g,b) which appear in Fig.3 (without the quadrupole) so the gluon,antigluon linked trefoil give us the $SU(3)_c$ color symmetry underlining QCD as described by Griffiths [9],Section 9.1.For example when another quark is added after a rotation ω a red gluon may unite with an anti-blue to build r anti-b, then a following rotation would bring r to anti-r ,and so on before flowing down to P^3 In this way we can find 9 gluon pairs r anti-r,r anti- b ,r anti- g;b anti-r,b anti-b,b anti- g;g anti-r,g anti-b,g anti- g that are a basis for $SU(3)_c$ symmetry.

Finally Adams [1] p 273 also envisages the 3 colors r,b,g as three extra dimensions in a 6-space

Acknowledgement

This contribution is a revision of a paper already submitted to PMSA 178X that was unfortunately published without the Figures

Fig.1 The Coxeter Polytope

Fig.2 Cayley Surface in Elliptic Space

Fig.3 Interior of Cayley Surface

4 Bibliography

References

- [1] C.C. Adams, The Knot Book, Freeman, 1994.
- [2] J.S.Arvin, Knots on a Torus: A Model of Elementary Particles, symmetry,4(2013)39-115.
- [3] W.Bruce and I.Miyamoto, Abelian Surfaces of Type (1,3) and Quartic Surfaces, J. Algebraic Geometry 3(1994)173-222.Benjamin Brau, Determining the Muon Mass in an Instructional Laboratory, Am.J.Phys.73(2010)64-70;arXiv:0907.5641.
- [4] Benjamin Brau, Determining the Muon Mass in an Instructional Laboratory, Am.J.Phys. 73(2010)64-70;arXiv:0907.5641.
- [5] H.S.M.Coxeter, The Polytope 221 whose 27 Vertices Correspond to the Lines on the General Cubic Surface, Am. Mathematical Soc.62(1940)467-486.
- [6] H.S.M.Coxeter, Regular Complex Polytopes, Camb.Univ.Press, Second Edition(1991).
- [7] J.A.de Wet, Icosahedral Symmetry in the MSSM, International Mathematical Forum 5(2010) 291-300. Available on-line under Hikari,Ltd.
- [8] M.B.Green,J.H.Schwarz and E.Witten, Superstring Theory ,Camb.Univ.Press (1998).
- [9] David Griffiths, Introduction to Elementary Particles, WILEY-VCH(2004).
- [10] H.Kedia, I.Bialynicki, D.Peralta-Salas and William Irvine, Tying Knots in LightFields,arXiv:1302.0342.

-
- [11] Bruce Hunt, The Geometry of Some Arithmetic Quotients, Lecture Notes in mathematics, 1637, Springer(1996).
- [12] Lei-Yang, Hessian Polyhedra Invariant Theory and Appell Hypergeometric Partial Differential Equations, arXiv:math/0412065v2.
- [13] <http://members.home.nl/fg.marcelis/ellip-cayley.html>.
- [14] Kunio Murasugi, Knot Theory and its Applications, Birkhauser, Boston(1996).
- [15] Carlo Rovelli, Loop Quantum Gravity, arXiv:gr-qc/971008.
- [16] R. Slansky, Group Theory for Unified Model Building, Reprinted in 'Unity of Forces in the Universe' Ed. A. Lee, World Scientific(1997).
- [17] Sundance O. Billson Thompson, Fotini Markopoulos and Lee Smolin, Quantum Gravity and the Standard Model, On-Line (2014)
- [18] Westy, Platonic Tessellations of Riemann Surfaces, <http://westy31.xs4all.nl/Geometry/Geometry.html>.

RETRACTED