ON DEGREE OF APPROXIMATION BY PRODUCT MEANS OF CONJUGATE SERIES OF A FOURIER SERIES

S. K. Paikray¹, R. K. Jati², N. C. Sahoo³, U.K.Misra⁴

¹P.G.Department of Mathematics, Ravenshaw University, Cuttack-753003, Odisha, India

²Department of Mathematics, DRIEMS, Tangi, Cuttack, Odisha, India

³Department of Mathematics, S.B. Women's College(Auto), Cuttack, Odisha, India

⁴Science and Technology, Pallur Hills, Golanthara-761008, Odisha, India

Keywords: Degree of Approximation $f \in Lip(\alpha, r)$ class of function, (E, q) mean, $(\overline{N}, p_n)$ mean, $(E, q)(\overline{N}, p_n)$ product mean, Fourier series, Conjugate of the Fourier series, Lebesgue integral

Abstract: In this paper a theorem on degree of approximation of a function $f \in Lip(\alpha, r)$ by product summability $(E, q)(\overline{N}, p_n)$ of conjugate series of Fourier series associated with f has been established.

1. Introduction

Let $\sum a_n$ be a given infinite series with the sequence of partial sums $\{s_n\}$. Let $\{p_n\}$ be a sequence of positive real numbers such that

$$P_n = \sum_{v=0}^n p_v \rightarrow \infty, \quad \text{as } n \rightarrow \infty, \quad (P_{-i} = p_{-i} = 0, i \geq 0). \quad (1.1)$$

The sequence –to–sequence transformation

$$t_n = \frac{1}{P_n} \sum_{v=0}^n p_v s_v \quad (1.2)$$

defines the sequence $\{t_n\}$ of the $(\overline{N}, p_n)$ -mean of the sequence $\{s_n\}$ generated by the sequence of coefficient $\{p_n\}$. If

$$t_n \rightarrow s, \quad \text{as } n \rightarrow \infty \quad (1.3)$$

then the series $\sum a_n$ is said to be $(\overline{N}, p_n)$ summable to s .

The conditions for regularity of $(\overline{N}, p_n)$ -summability are easily seen to be [1]

$$\begin{cases} (i) P_n \rightarrow \infty, \text{ as } n \rightarrow \infty. \\ (ii) \sum_{i=0}^n p_i \leq C |P_n|, \text{ as } n \rightarrow \infty. \end{cases} \quad (1.4)$$

The sequence –to–sequence transformation, [1]

$$T_n = \frac{1}{(1+q)^n} \sum_{v=0}^n \binom{n}{v} q^{n-v} s_v \quad (1.5)$$

defines the sequence $\{T_n\}$ of the (E, q) mean of the sequence $\{s_n\}$.

If

$$T_n \rightarrow s, \quad \text{as } n \rightarrow \infty, \quad (1.6)$$

then the series $\sum a_n$ is said to be (E, q) summable to s .

Clearly (E, q) method is regular. Further, the (E, q) transform of the (N, p_n) transform of $\{s_n\}$ is defined by

$$\begin{aligned}\tau_n &= \frac{1}{(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} t_k \\ &= \frac{1}{(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{v=0}^k P_v s_v \right\}\end{aligned}\quad (1.7)$$

If

$$\tau_n \rightarrow s, \text{ as } n \rightarrow \infty, \quad (1.8)$$

then $\sum a_n$ is said to be $(E, q)(N, p_n)$ -summable to s .

Let $f(t)$ be a periodic function with period 2π and L -integrable over $(-\pi, \pi)$. The Fourier series associated with f at any point x is defined by

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) \equiv \sum_{n=0}^{\infty} A_n(x) \quad (1.9)$$

and the conjugate series of the Fourier series (1.9) is

$$\sum_{n=1}^{\infty} (b_n \cos nx - a_n \sin nx) \equiv \sum_{n=1}^{\infty} B_n(x) \quad (1.10)$$

Let $S_n(f; x)$ be the n -th partial sum of (1.10). The L^∞ -norm of a function $f: R \rightarrow R$ is defined by

$$\|f\|_\infty = \sup \{ |f(x)| : x \in R \} \quad (1.11)$$

and the L_v -norm is defined by

$$\|f\|_v = \left(\int_0^{2\pi} |f(x)|^v dx \right)^{\frac{1}{v}}, \quad v \geq 1 \quad (1.12)$$

The degree of approximation of a function $f: R \rightarrow R$ by a trigonometric polynomial $P_n(x)$ of degree n under norm $\|\cdot\|_\infty$ defined by [5].

$$\|P_n - f\|_\infty = \sup \{ |P_n(x) - f(x)| : x \in R \} \quad (1.13)$$

and the degree of approximation $E_n(J)$ of a function $J \in L_v$ is given by

$$E_n(f) = \min_{P_n} \|P_n - f\|_v \quad (1.14)$$

A function f is said to satisfy Lipschitz condition (here after we write $f \in Lip_\alpha$) if

$$|f(x+t) - f(x)| = O(|t|^\alpha), \quad 0 < \alpha \leq 1 \quad (1.15)$$

and $f(x) \in Lip(a, r)$, for $0 \leq x \leq 2\pi$, if

$$\left(\int_0^{2\pi} |f(x+t) - f(x)|^r dx \right)^{\frac{1}{r}} = O(|t|^\alpha), \quad 0 < \alpha \leq 1, r \geq 1, t > 0 \quad (1.16)$$

We use the following notation throughout this paper :

$$\psi(t) = \frac{1}{2} \{f(x+t) - f(x-t)\}, \quad (1.17)$$

and

$$\overline{K}_n(t) = \frac{1}{\pi(1+q)^n} \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{v=0}^k p_v \frac{\cos \frac{t}{2} - \cos \left(v + \frac{1}{2} \right) t}{\sin \frac{t}{2}} \right\} \quad (1.18)$$

Further, the method $(E, q)(N, p_n)$ is assumed to be regular .

2. Known Theorem

Dealing with The degree of approximation by the product $(E, q)(C, 1)$ -mean of Fourier series, Nigam et al [3] proved the following theorem:

Theorem 2.1. If a function f is 2π -periodic and belonging to class $\underline{Lipa}$, then its degree of approximation by $(E, q)(C, 1)$ summability mean on its Fourier series $\sum_{n=0}^{\infty} A_n(t)$ is given by

$$\|E_n^q C_n^1 - f\|_{\infty} = O\left(\frac{1}{(n+1)^{\alpha}}\right), 0 < \alpha < 1$$

where $E_n^q C_n^1$ represents the (E, q) transform of $(C, 1)$ transform of $S_n(f; x)$.

Recently, Misra et al [2] proved the following theorem using $(E, q)(N, p_n)$ mean of conjugate series of the Fourier series :

Theorem 2.2. If f be 2π -Periodic function of class $\underline{Lipa}$, then degree of approximation by the product $(E, q)(\overline{N}, p_n)$ summability means of the conjugate series (1.10) of Fourier series (1.9) is

$$\|\tau_n - f\|_{\infty} = O\left(\frac{1}{(n+1)^{\alpha}}\right), 0 < \alpha < 1$$

given where τ_n is as defined in 1.7).

3. Main theorem

In this paper, we have proved a theorem on degree of approximation by the product mean $(E, q)(N, p)$ of conjugate series of Fourier series of a function of class $Lip(a, r)$. We prove:

Theorem.3.1. If f be 2π -Periodic function of class $\underline{Lipa}$, then degree of approximation by the product $(E, q)(\overline{N}, p_n)$ summability means of the conjugate series of Fourier series (1.10) is given

$$\|\tau_n - f\|_{\infty} = O\left(\frac{1}{(n+1)^{\alpha-1/r}}\right), 0 < \alpha < 1$$

by where τ is as defined in (1.7).

4. Required Lemmas

We require the following Lemmas to prove the theorem.

Lemma 4.1

$$|\overline{K}_n(t)| = O(n), 0 \leq t \leq \frac{1}{n+1}$$

Proof.

For $0 \leq t \leq \frac{1}{n+1}$ we have $\sin nt \leq n \sin t$ then

$$\begin{aligned}
 |\overline{K}_n(t)| &= \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \frac{\cos \frac{t}{2} - \cos \left(\nu + \frac{1}{2} \right) t}{\sin \frac{t}{2}} \right\} \right| \\
 &\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \frac{\cos \frac{t}{2} - \cos \nu t \cdot \cos \frac{t}{2} + \sin \nu t \cdot \sin \frac{t}{2}}{\sin \frac{t}{2}} \right\} \right| \\
 &\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \left(\frac{\cos \frac{t}{2} \left(2 \sin^2 \nu \frac{t}{2} \right)}{\sin \frac{t}{2}} + \sin \nu t \right) \right\} \right| \\
 &\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \left(O \left(2 \sin \nu \frac{t}{2} \sin \nu \frac{t}{2} \right) + \nu \sin t \right) \right\} \right| \\
 &\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu (O(\nu) + O(\nu)) \right\} \right| \\
 &\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \frac{O(k!)}{P_k} \sum_{\nu=0}^k p_\nu \right| \\
 &= O(n)
 \end{aligned}$$

This proves the lemma.

Lemma 4.2

$$|\overline{K}_n(t)| = O\left(\frac{1}{t}\right), \text{ for } \frac{1}{n+1} \leq t \leq \pi$$

Proof.

For $\frac{1}{n+1} \leq t \leq \pi$, by Jordan's lemma, we have $\sin\left(\frac{t}{2}\right) \geq \frac{t}{\pi}$

Then

$$\begin{aligned}
 |\overline{K}_n(t)| &= \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \frac{\cos \frac{t}{2} - \cos \left(\nu + \frac{1}{2} \right) t}{\sin \frac{t}{2}} \right\} \right| \\
 &= \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \frac{\cos \frac{t}{2} - \cos \nu \frac{t}{2} \cdot \cos \frac{t}{2} + \sin \nu \frac{t}{2} \cdot \sin \frac{t}{2}}{\sin \frac{t}{2}} \right\} \right|
 \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{\pi(1+q)^n} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k \frac{\pi}{2t} p_\nu \left(\cos \frac{t}{2} \left(2 \sin^2 \nu \frac{t}{2} \right) + \sin \nu \frac{t}{2} \cdot \sin \frac{t}{2} \right) \right\} \right| \\
&\leq \frac{\pi}{2\pi(1+q)^n t} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \right\} \right| = \frac{1}{2(1+q)^n t} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \right\} \right| \\
&= \frac{1}{2(1+q)^n t} \left| \sum_{k=0}^n \binom{n}{k} q^{n-k} \right| \\
&= O\left(\frac{1}{t}\right).
\end{aligned}$$

This proves the lemma.

5. Proof of theorem 3.1

Using Riemann –Lebesgue theorem, we have for the n -th partial sum $s_n(\bar{f}; x)$ of the conjugate Fourier series (1.10) of $f(x)$, following Titchmarsh [4]

$$\overline{s_n}(f; x) - f(x) = \frac{2}{\pi} \int_0^\pi \psi(t) \overline{K_n} dt$$

the (N, pn) transform of $\overline{s_n}(f; x)$ using (1.2) is given by

$$t_n - f(x) = \frac{2}{\pi P_n} \int_0^\pi \psi(t) \sum_{k=0}^n p_k \frac{\cos \frac{t}{2} - \sin\left(n + \frac{1}{2}\right) t}{2 \sin\left(\frac{t}{2}\right)} dt$$

denoting the $(E, q)(N, pn)$ transform of $\overline{s_n}(f; x)$ by τ_n , we have

$$\|\tau_n - f\| = \frac{2}{\pi(1+q)^n} \int_0^\pi \psi(t) \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{\nu=0}^k p_\nu \frac{\cos \frac{t}{2} - \sin\left(\nu + \frac{1}{2}\right) t}{2 \sin\left(\frac{t}{2}\right)} \right\} dt$$

$$= \int_0^\pi \psi(t) \overline{K_n}(t) dt$$

$$= \left\{ \int_0^{\frac{1}{n+1}} + \int_{\frac{1}{n+1}}^\pi \right\} \psi(t) \overline{K_n}(t) dt$$

$$= I_1 + I_2, \text{ say}$$

$$(5.1)$$

Now

$$\begin{aligned}
 |I_1| &= \frac{2}{\pi(1+q)^n} \left| \int_0^{1/n+1} \psi(t) \sum_{k=0}^n \binom{n}{k} q^{n-k} \left\{ \frac{1}{P_k} \sum_{v=0}^k p_v \frac{\cos \frac{t}{2} - \cos \left(v + \frac{1}{2} \right) t}{2 \sin \frac{t}{2}} \right\} dt \right| \\
 &\leq \left| \int_0^{1/n+1} \psi(t) \overline{K}_n(t) dt \right| \\
 &= \left(\int_0^{1/n+1} (\psi(t))^r dt \right)^{\frac{1}{r}} \left(\int_0^{1/n+1} (\overline{K}_n(t))^s dt \right)^{\frac{1}{s}}, \text{ using Holder's inequality} \\
 &\leq O\left(\frac{1}{(n+1)^\alpha}\right) \left(\int_0^{1/n+1} n^s dt \right)^{\frac{1}{s}} \quad \text{using Lemma 4.1} \\
 &= O\left(\frac{1}{(n+1)^\alpha}\right) \left(\frac{n^s}{s+1} \right)^{\frac{1}{s}} \\
 &= O\left(\frac{n}{(n+1)^{\alpha+\frac{1}{s}}}\right). \\
 &= O\left(\frac{1}{(n+1)^{\alpha+\frac{1}{s}-1}}\right) = O\left(\frac{1}{(n+1)^{\alpha-\frac{1}{r}}}\right) \tag{5.2}
 \end{aligned}$$

Next

$$\begin{aligned}
 |I_2| &\leq \left(\int_{\frac{1}{n+1}}^{\pi} (\psi(t))^r dt \right)^{\frac{1}{r}} \left(\int_{\frac{1}{n+1}}^{\pi} (\overline{K}_n(t))^s dt \right)^{\frac{1}{s}} \quad \text{using Holder's inequality} \\
 &\leq O\left(\frac{1}{(n+1)^\alpha}\right) \left(\int_{\frac{1}{n+1}}^{\pi} \left(\frac{1}{t}\right)^s dt \right)^{\frac{1}{s}} \quad \text{using Lemma 4.2} \\
 &= O\left(\frac{1}{(n+1)^\alpha}\right) \left(\frac{1}{n+1} \right)^{\frac{1-s}{s}} \\
 &= O\left(\frac{1}{(n+1)^{\alpha-\frac{1}{r}}}\right) \tag{5.3}
 \end{aligned}$$

Then from (5.2) and (5.3) , we have

$$|\tau_n - f(x)| = O\left(\frac{1}{(n+1)^{\alpha - \frac{1}{r}}}\right), \text{ for } 0 < \alpha < 1, r \geq 1.$$

Hence,

$$\|\tau_n - f(x)\|_{\infty} = \sup_{-\pi < x < \pi} |\tau_n - f(x)| = O\left(\frac{1}{(n+1)^{\alpha - \frac{1}{r}}}\right), 0 < \alpha < 1, r \geq 1$$

This completes the proof of the theorem.

References

- [1] Hardy, G.H: Divergent series, First edition, Oxford University press 70(19).
- [2] Misra, U.K., Misra, M., Padhy, B.P. and Buxi, S.K: On degree of approximation by product means of conjugate series of Fourier series", International Jour. of Math. Sciences, and Engineering Applications, ISSN 0973 – 9424, Vol.6 No.122 (Jan. 2012), pp 363 – 370.
- [3] Nigam, H.K and Ajay Sharma: On degree of Approximation by product means, Ultra Scientist of Physical Sciences, Vol.22 (3) M, 889-894, (2010).
- [4] Titchmarsh, E.C: The theory of functions, oxford university press, p.p402-403(1939).
- [5] Zggmund , A : Trigonometric Series , second Edition ,Vol.I , Cambridge University press , Cambridge , (1959).