An Elegant Proof that the Catalan's Constant is Irrational

Edigles Guedes¹ and Prof. Dr. Raja Rama Gandhi²

Number Theorsit, Brazil¹

Resorce person in Math for Oxford University Press and Professor at BITS-Vizag²

ABSTRACT. We use the contradiction method for prove that the Catalan's constant is irrational.

1. INTRODUCTION

In Mathematics, the Catalan's constant [1] is defined by

$$G := \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2}. \quad (1.1)$$

It is not known whenever G is irrational, let alone transcendental. The first open problem is the subject of our paper.

The Catalan's constant was named after Eugène Charles Catalan (20 May 1814 – 14 February 1894), a French and Belgian mathematician.

2. THE PROOF

THEOREM. *The Catalan's constant is irrational.*

Proof. We will use the *reductio ad absurdum*.

By hypothesis, we suppose that G is a rational number. Of course, there exist two positive integers a and b , such that $G = a/b$, where, clearly, $b > 1$. Firstly, we define the number

$$x := \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \cdot \left| \sum_{n=0}^b \frac{(-1)^n}{(2n+1)^2} \right|. \quad (2.1)$$

If G is rational, then x is an integer. We substitute $G = a/b$ into this definition to find

$$x = \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \cdot \left| \frac{a}{b} - \sum_{n=0}^b \frac{(-1)^n}{(2n+1)^2} \right| = \left| \frac{a(2b+1)!^2}{4^b (b!)^2} - \sum_{n=0}^b \frac{(-1)^n (2b+1)!^2}{(2n+1)^2 4^b b((b-1)!)^2} \right|. \quad (2.2)$$

It is obvious that the first term is an integer; because, for $b > 1$, then $4^b (b!)^2 < (2b+1)!^2$. The second term is an integer; because, for $b > 1$, then $(2n+1)^2 4^b b((b-1)!)^2 < (2b+1)!^2$. Hence x is an integer.

We, now, demonstrate that $0 < x < 1$.

First, to demonstrate that x is strictly positive, we insert the series representation of G into the definition of x and we find

$$\begin{aligned} x &= \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} - \sum_{n=0}^b \frac{(-1)^n}{(2n+1)^2} \right| = \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{(-1)^n}{(2n+1)^2} \right| = \\ &= \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{\cos(\pi n)}{(2n+1)^2} \right| > \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \int_{b+1}^{\infty} \frac{\cos(\pi x)}{(2x+1)^2} dx \right| = \\ &= \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| -\frac{1}{4} \pi \operatorname{Ci} \left(\left(b + \frac{3}{2} \right) \pi \right) - \frac{\cos(\pi b)}{4b+6} \right| > 0. \end{aligned} \quad (2.3)$$

On the other hand, for all terms with $2n+1 \geq 2b+2$, i.e., $2n \geq 2b+1$, we have the upper estimate

$$\frac{(2b+1)!}{(2n+1)!} \leq \frac{1}{(2b+2)^{2n-2b}}. \quad (2.4)$$

This inequality is strict for every $2n + 1 \geq 2b + 3$, i.e., $n \geq b + 1$. Thereof, we substitute (1.1) and (2.4) in (2.1)

$$\begin{aligned}
 x &= \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} - \sum_{n=0}^b \frac{(-1)^n}{(2n+1)^2} \right| \\
 &= \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{(-1)^n}{(2n+1)^2} \right| < \frac{(2b+1)!^2}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{(-1)^n}{(2n+1)!^2} \right| \\
 &= \frac{1}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{(-1)^n (2b+1)!^2}{(2n+1)!^2} \right| < \frac{1}{4^b b((b-1)!)^2} \left| \sum_{n=b+1}^{\infty} \frac{(-1)^n}{(2b+2)^2} \right| \\
 &= \frac{1}{4^b b((b-1)!)^2} \left| -\frac{(-1)^b}{4b^2+8b+5} \right| < 1. \tag{2.5}
 \end{aligned}$$

Since there is no integer strictly between 0 and 1, we have get in contradiction, and so G must be irrational. $\square$

REFERENCES

- [1] http://en.wikipedia.org/wiki/Catalan's_constant, available in August 9, 2013.