

Insect antifeedant potent halogen substituted phenyl chalcones

R. Sundararajan¹, R. Arulkumar¹, S. Vijayakumar¹, D. Kamalakkannan¹,
R. Suresh¹, S. John Joseph¹, K. Ranganathan¹, S. P. Sakthinathan¹,
G. Vanangamudi¹, G. Thirunarayanan^{2,*}

¹ PG & Research Department of Chemistry, Government Arts College, C-Mutlur,
Chidambaram - 608102, India.

² Department of Chemistry, Annamalai University, Annamalainagar - 608002, Inida.

*E-mail address: drgtnarayanan@gmail.com

ABSTRACT

Some 2',3',4'-trichlorophenyl chalcones [(*E*)-1-(2,3,4-trichlorophenyl)-3-(substituted phenyl)-2-propen-1-ones] have been synthesised using sulfated Titania catalyzed solvent-free aldol condensation between 2,3,4-trichloroacetophenone and substituted benzaldehydes. The purities of synthesised chalcones were checked by their analytical, physical and spectroscopic data reported in literature. The insect antifeedant activities of these chalcones have been studied using 4th instar larvae *Achoea Janata L* by castor leaf disc bio-assay method. The chloro substituted chalcones shows significant insect antifeedant activity.

Keywords: Substituted styryl 2',3',4'-trichlorophenyl ketones; sulfated Titania; IR and NMR spectra; Insect antifeedant activities

1. INTRODUCTION

Chalcones are α , β -unsaturated ketones. In their structure, they contain $-\text{CO}-\text{CH}=\text{CH}-$ moiety between alkyl, alkyl or aryl, aryl or alkyl aryl groups with or without substituents. Numerous chalcone derivatives are used as agrochemicals [1] and drugs [2] due to their structurally important multipronged activities by the presence of carbonyl, vinyl and polar substituents in their structure. The important multipronged activities of chalcones are anti-microbial [3], anti-cancer [4], anti-viral [5], anti-malarial [6], anti-oxidant [7], anti-proliferative [8], anti-AIDS [9], anti-tuberculosis [10], anti-plasmodial [11], anti-inflammatory [12], anti-leishmanial [13], anti-convulsant [14], cytotoxic [15], anti-parasitic [16], anti-angiogenic [17], anti-mycobacterial [13], anti-trichomonal [19], anti-amastigote [20], radio protective [21], and insect antifeedant activities [22].

These chalcones also used as corrosion inhibitors in iron and steel utensil and equipment manufacturing industries [23].

These chalcones are also precursor key intermediate for synthesis of important biologically active higher organic heterocycles such as flavones, flavonoids, chromones, auronones andazole related compounds [24,25].

In agricultural field they play with much more important role as insect antifeedants. Dasharathi et al., have studied the solvent assisted synthesis and insect antifeedant activities of some multi-substituted naphthyl styryl ketones [26]. Vibhut et al., have studied the insect antifeedant activities of some chalcones by invitro bio-assay method [27].

Thirunarayanan and his co-workers studied the synthesis and insect antifeedant activities of halogen substituted 1 and 2-naphthyl styryl ketones [28], heterocyclic ketones [29], halo substituted phenyl styryl ketones [30], ω -bromo acyl compounds [31] and halo epoxides [32]. With the above view, there is no report available for the synthesis of chalcones by perchloric acid catalyzed condensation and the study of insect antifeedant activities of 2',3',4'-trichlorophenyl chalcones in literature in the past.

Therefore, the authors taken efforts for studying the insect antifeedant activities of the above chalcones by solvent-free synthesis and checked the purities by reported analytical, physical constants and spectral data [37-41].

2. EXPERIMENTAL

2.1. General

All chemicals used were procured from Sigma-Aldrich and Merck chemical companies. Mettler FP51 melting point apparatus was used for determining the melting point of all synthesized chalcones in open glass capillaries and are uncorrected.

The absorption maxima (λ_{\max} , nm), of all chalcones were determined by recording the ultraviolet spectra on ELICO BL 222 ultraviolet spectrophotometer using spectral grade methanol. The AVATAR-300 Fourier transform spectrophotometer was used for recording infrared spectra (KBr, 4000-400 cm^{-1}) of all chalcones in KBr disc.


The Bruker AV400 series type NMR spectrometer was utilized for recording NMR spectra of all chalcones, operating at 400MHz for ^1H and 100 MHz for ^{13}C spectra in CDCl_3 solvent using TMS as internal standard. Mass spectra of all synthesised chalcones were recorded on SHIMADZU mass spectrometer using chemical ionization technique.

2.2. Synthesis of substituted styryl 2', 3', 4'-trichlorophenyl ketones

Appropriate equi-molar quantities of 2',3',4'-trichloroacetophenone (2 mmol) and various substituted benzaldehydes (2 mmol) and 0.15 g of sulfated titania [33,34] was taken in a 50 mL beaker, closed with the lid.

This mixture was subjected to microwave irradiation for 2-4 minutes at 650W (Scheme 1) (Samsung, Microwave Oven, 100-700W). After completion of the reaction, dichloromethane (20 mL) was added, followed by simple filtration.

The solution was concentrated and purified by re-crystallization. The synthesized chalcones were characterized by their physical constants, UV, IR, ^1H and ^{13}C NMR and Mass spectral data. Analytical and Mass spectral data are presented in Table 1.



Where X = H, 3-Br, 4-Br, 3-Cl, 4-Cl, 2-OCH₃, 4-OCH₃, 2-CH₃, 4-CH₃, 3-NO₂, 4-NO₂

Scheme 1. Synthesis of 2',3',4'-trichlorophenyl chalcones.

Table 1. Physical constants, analytical and Mass fragments(m/z) of substituted styryl 2',3',4'-trichlorophenyl ketones.

Entry	X	M. F.	M. W.	m.p.(°C)	Mass (m/z)
1	H	C ₁₅ H ₉ OCl ₃	310	161-162	310[M ⁺]
2	3-Br	C ₁₅ H ₈ OCl ₃ Br	387	148-149	387 [M ⁺], 395[M ⁸⁺], 393[M ⁶⁺], 391[M ⁴⁺], 389[M ²⁺]
3	4-Br	C ₁₅ H ₈ OCl ₃ Br	387	160-161	387 [M ⁺], 395[M ⁸⁺], 393[M ⁶⁺], 391[M ⁴⁺], 389[M ²⁺]
4	3-Cl	C ₁₅ H ₈ OCl ₄	344	138-139	344[M ⁺], 352[M ⁸⁺], 350[M ⁶⁺], 348[M ⁶⁺], 346[M ²⁺]
5	4-Cl	C ₁₅ H ₈ OCl ₄	344	165-166 (162–165)[35]	344[M ⁺], 352[M ⁸⁺], 350[M ⁶⁺], 348[M ⁶⁺], 346[M ²⁺]
6	2-OCH ₃	C ₁₆ H ₁₁ O ₂ Cl ₃	340	116-117	340[M ⁺], 346[M ⁶⁺], 344[M ⁴⁺], 342[M ²⁺]
7	4-OCH ₃	C ₁₆ H ₁₁ O ₂ Cl ₃	340	127-128	340[M ⁺], 346[M ⁶⁺], 344[M ⁴⁺], 342[M ²⁺]
8	2-CH ₃	C ₁₆ H ₁₁ OCl ₃	324	101-102	324[M ⁺], 330[M ⁶⁺], 328[M ⁴⁺], 326[M ²⁺]
9	4-CH ₃	C ₁₆ H ₁₁ OCl ₃	324	98-99	324[M ⁺], 330[M ⁶⁺], 328[M ⁴⁺], 326[M ²⁺]
10	3-NO ₂	C ₁₅ H ₈ O ₃ Cl ₃ N	355	188-189	355[M ⁺], 359[M ⁴⁺], 357[M ²⁺]
11	4-NO ₂	C ₁₅ H ₈ O ₃ Cl ₃ N	355	190-191	355[M ⁺], 359[M ⁴⁺], 357[M ²⁺]

2. 3. Measurement of insect antifeedant activities of 2',3',4'-trichlorophenyl chalcones

The castor Leaf discs of a diameter of 1.85 cm were punched with the petioles intact. All chalcones were dissolved in acetone at a concentration of 200 ppm dipped for 5 minutes. The air-dried leaf discs were placed in 1 L beaker containing little water in order to facilitate translocation of water. Therefore the leaf discs remains fresh throughout the duration of the rest, 4th instar larvae of the test insect, which had been preserved on the leaf discs of all

3. RESULTS AND DISCUSSION

The multipronged activities present in different 2',3',4'-trichlorophenyl chalcones are intended to examine their insect antifeedant activities against castor *semilooper*. The larvae of *Achoea janata* L were reared as described on the leaves of castor *Ricinus communis* in the laboratory at the temperature range of 26° C ±1 °C and a relative humidity of 75-85 %. The leaf – disc bioassay method was used against the 4th instar larvae to measure the antifeedant activity. The 4th instar larvae were selected for testing because the larvae at this stage feed very voraciously.

The results of the antifeedant activity of substituted styryl 2',3',4'-trichlorophenyl ketones were presented in Table 2 and it reveals that the compounds 2-5 were found to reflect remarkable antifeedant among all other naphthones. This test is performed with the insects which are only two-leaf disc soaked under the solution of this compound. Compounds 2-5 also show enough antifeedant activity but lesser than 5. Further compound 5 was subjected to measure the antifeedant activity at different 50, 100, 150 ppm concentrations and the observation reveals that as the concentrations decreased, the activity also decreased. It is observed from the results in Table 3 and that the 4-chlorostyryl 2',3',4'-trichlorophenyl ketone 5 showed an appreciable antifeedant activity at 150 ppm concentration.

4. CONCLUSIONS

Totally eleven substituted styryl 2',3',4'-trichlorophenyl ketones [(*E*)-1-(2,3,4-trichlorophenyl)-3-(substituted phenyl)-2-propen-1-ones] were synthesised using sulfated Titania catalyzed solvent-free aldol condensation. The purities of synthesised chalcones were characterized by their analytical, physical and spectroscopic data. The insect antifeedant activities of these chalcones have been studied using 4th instar larvae *Achoea Janata L* by castor leaf disc bio-assay method. The chloro substituted chalcones(5) 4-chlorostyryl 2', 3', 4'-trichlorophenyl ketone showed significant insect antifeedant activity.

Acknowledgement

The authors thank to the DST NMR facility unit, Department of Chemistry, Annamalai University, Annamalinagar - 608 002, for recording NMR spectra of all compounds.

References

- [1] Mirinda C. L., Aponso G. L., Stevens J. F., *Pure Appl Chem* 73 (2000) 1197-1208.
- [2] Monostory K., Tamasi V., Vereckey L., Perjesi P., *Toxicol* 184 (2003) 203-210.
- [3] Subramanian M., Vanangamudi G., Thirunarayanan G., *Spectrochim Acta*. 110A (2013) 116-123.
- [4] Modzelewska A., Pettit C., Achanta G., Davidson N. E., Huang P., Khan R. K., *Bioorg Med Chem* 14 (2006) 3491-3495.
- [5] Ahmad A. L., Dowsett A. B., Tyrrell D. A., *Antiviral Res.* 8 (1987) 27-39.
- [6] Liu M., Wilairat M. L., Croft, S. L., Tan A. L. C., Go M., *Bioorg Med Chem.* 11 (2003) 2729-2738.

-
- [7] Weber W. M., Hunsaker L. A., Abcouwer S. F., Deck L. M., Jagt D. L. V., *Bioorg Med Chem.* 13 (2005) 3811-3820.
- [8] Liu X., Go M. L., *Bioorg Med Chem.* 14 (2006) 153-163.
- [9] Deng J., Sanchez T., Lalith Q. A. M., Dayam R., Yunes R. A., Garofalo A., et al. *Bioorg Med Chem.* 15 (2007) 4985-5002.
- [10] Lin Y. M., Zhou Y., Flavin M. T., Zhou L. M., Nie W., Che F. C., *Bioorg Med Chem.* 10 (2002) 2795-2802
- [11] Arulkumaran R., Sundararajan R., Vanangamudi G., Subramanian M., Ravi K., Sathiyendiran V. et al.. *Iup J Chem.* 3 (2010) 82-98.
- [12] Kotra V., Ganapathy S., Adapa S. R., *Indian J Chem* 49B (2010) 1109-1116.
- [13] Reichwald C., Shimony O., Sierra N. S., Jaffe C. L., Kunick C., *J Med Chem.* 51 (2008) 659-665.
- [14] Said M. M., Ahmed A. A. E., Alfay A. T. E., *Arch Pharm Res.* 27 (2004) 1194-1201.
- [15] Bhat B. A., Dhar K. L., Puri A. K., Saxena A. K., Shanmugavel M., Quazi G. N., *Bioorg Med.Chem.* 15 (2005) 3177-3180.
- [16] Tragar W., Jenson J. B., *Science.* 193 (1976) 673-675.
- [17] Nem N. H., Kim. Y., You Y. J., Hong D. H., Kim H. M., Ahn B. Z., *Eur J Med Chem.* 38(2) (2003) 179-187.
- [18] Pushkar N. K., Balawant S. J., *Alternative Medicine: Herbal Drugs and their Critical appraisal - Part II*”, *Progress in Drug Research.* 2001; 57: (E. Jucker, Ed.) Birkhäuser Verlag, Basel (Switzerland).
- [19] Oyedapo A. O., Mankanju V. O., Adewunmi C. O., Iwalewa E. O., Adenowo T. K., *Afr J Trad CAM.* 1 (2004) 55-62.
- [20] Suryawashni S. N., Chandra N., Kumar P., Porwal J., Gupta S., *Eur J Med Chem.* 43 (2008) 2473-2478.
- [21] Blickenstaff R.T., Hanson W. R., Reddy S., Witt R., *Bioorg Med Chem.* 3 (1995) 917-922.
- [22] Thirunarayanan G., Surya S., Srinivasan, S., Vanangamudi G., Sathiyendiran V., *Spectrochim Acta.* 75A (2010) 152-156.
- [23] Fouda A. S., Shafie A. A. E., Gadow H. S., *Portugaliae Electrochim Acta.* 20 (2002) 13-23.
- [24] Sekar K. G., Thirunarayanan G., *Int J Sci Res Know.* 1 (2013) 299-307.
- [25] Thirunarayanan G., *Asian J Chem.* 15 (2003) 907-910.
- [26] Dasharathi D., Netaji R., Basheer M. A., Vibhute Y. B., *Ultra Science.* 17 (2005) 89-92.
- [27] Nalwar Y. S., Sayyed M. A., Mokle S. S., Zanwar P, R., Vibhute Y. B., *World J. Chem.* 4(2009)123-126.
- [28] Thirunarayanan G., *J Indian Chem Soc.* 85(2008)447-451.
- [29] Janaki P., Sekar K. G., Thirunarayanan G., *J Saudi Chem Soc.* (2012). Doi:org.10.1016/j.jscs.2012.11.013.

- [30] Thirunarayanan G., Thirumurthy K., Vanangamudi G., Subramanian M., Arulkumaran R., Kamalakkannan D. et al. *Elixir Org Chem.* 45 (2012) 7898-7905.
- [31] Thirunarayanan G., *J Saudi Chem Soc.* (2011). doi:10.1016/j.jscs.2011.12.003.
- [32] Thirunarayanan G., *J Saudi Chem Soc.* (2011). doi:10.1016/j.jscs.2011.10.011.
- [33] Krishnakumar B., Velmurugan R., Swaminathan M., *Catal Commun.* 12 (2011) 375-379.
- [34] Sundararajan R., Arulkumaran R., Vijayakumar S., Kamalakkannan D., Suresh R., John Joseph S., Ranganathan K., Vanangamudi G., Subramanian M., Thirunarayanan G., Muthuvel I., Krishnakumar B., *Q-Science Connect.* 2013. DOI: <http://dx.doi.org/10.5339/connect.2013.30>
- [35] Hoong-Kun F., Yeap C. S., Jagadeesh Prasad D., Nayak S. P., Laxmana K., *Acta Cryst Sec E* 67 (2011) o241.
- [36] Dethler V. G., *Chemical insect attractants and Repellants.* Blackistan, Philadeciphia, pp. 210. (1947).
- [37] R. Arulkumaran, S. Vijayakumar, R. Sundararajan, S. P. Sakthinathan, D. Kamalakkannan, R. Suresh, K. Ranganathan, G. Vanangamudi, G. Thirunarayanan, *International Letters of Chemistry, Physics and Astronomy* 4 (2012) 17-38.
- [38] K. Ranganathan, R. Suresh, D. Kamalakkannan, R. Arulkumaran, R. Sundararajan, S. P. Sakthinathan, S. Vijayakumar, G. Vanangamudi, K. Thirumurthy, P. Mayavel, G. Thirunarayanan, *International Letters of Chemistry, Physics and Astronomy* 4 (2012) 66-75.
- [39] R. Arulkumaran, S. Vijayakumar, R. Sundararajan, S. P. Sakthinathan, D. Kamalakkannan, R. Suresh, K. Ranganathan, P. R. Rajakumar, G. Vanangamudi, G. Thirunarayanan, *International Letters of Chemistry, Physics and Astronomy* 5 (2013) 21-38.
- [40] S. Vijayakumar, R. Arulkumaran, R. Sundararajan, S. P. Sakthinathan, R. Suresh, D. Kamalakkannan, K. Ranganathan, K. Sathiyamoorthy, V. Mala, G. Vanangamudi, G. Thirunarayanan, *International Letters of Chemistry, Physics and Astronomy* 9(1) (2013) 68-86.
- [41] Thirunarayanan G., Sekar K. G., *International Letters of Chemistry, Physics and Astronomy* 10 (2013) 18-34.

(Received 03 October 2013; accepted 07 October 2013)