

Determination of heavy metals of road deposited sediment in Ado-Ekiti, Nigeria using XRF Technique

O. T. Ogunmodede¹, O. O. Ajayi²

¹Department of Chemical Science, Afe Babalola University, Ado-Ekiti, Ekiti State, Nigeria

²Department of Industrial Chemistry, Federal University of Technology, Akure, Ondo State, Nigeria

ABSTRACT

In this work x-ray fluorescence(XRF) technology was used to evaluate the soil pollution with heavy metals (K, Ti, Cr, Mn, Fe, Cu, Zr) in rain run-off deposited metal sediment of road side soil in Ado Ekiti, Nigeria. The investigated sediment of road side was collected in open places along the road at different districts in Ado Ekiti. XRF was carried out at the laboratory of Obafemi Awolowo University centre for energy research using handheld thermo scientific energy-dispersive XRF analyzer. The experimental result indicate that the concentration of heavy elements in Adebayo road is the highest level detected while the road at new Iyin road is lowest and they are greater than the level detected in a control soil collected from a zone situated far from the road. For the majority of metals, pronounced maximum, concentrations were detected in the site. Anthropogenic releases give rise to highest concentrations of the metals relative to the normal background values and in some locations their levels exceed the alert level admitted by the Nigeria guideline.

Keywords: X –Ray fluorescence (XRF) technique; heavy metals; soil pollution; anthropogenic

1. INTRODUCTION

Sediments on road surfaces and in curbside areas are ubiquitous in urban and sub-urban drainage basins. These deposits are easily sampled, and provide a useful indicator of the degree of pollution status of a locate [1], curbside sediment and associated contaminants are typically available for mobilization and transportation to sub surface drainage system by storm water run-off. Established research has shown that sediments and dusts transported and stored in the urban environment have the potential to provide consider able loading of heavy metals to receiving water and water bodies particularly with changing environmental conditions [2]. Street sediments that accumulate along parameters in urban environments originate mainly from natural and anthropogenic sources. Heavy metal from natural sources vary significantly within catchment and may include materials transported by water from surrounding soils, pollutants from dry and net atmospheric deposition and biological materials from vegetation. Significant quality of particulate matter can also be attributed to anthropogenic sources such as abrasion of vehicular component and their exhaust emission, incinerators, power plants and foundry operations, type and road surface wear [3,4]. These deposits as street sediments have become an important medium for study of anthropogenic pollutants and their possible sources [5-9]. Urban street sediment has limited residence tunes and therefore provides a record of recent accumulation [10,11]. The attractiveness of non-destructive method and the ability to perform simultaneous multi-elemental determination has to an extensive application in industrial and research laboratories of accurate, precise and sensitive atomic and unclear analytical techniques for the investigation of different types of

materials. The main goal of the present research was to use XRF techniques in order to assess the heavy metals distribution in road side soil sediments in Ado- Ekiti Nigeria.

2. MATERIALS AND METHODS

Studied area was Ado Ekiti city which is at 7°31' N and 5°5' E. Fifty sediment samples were collected from ten roads from five districts of the city with the aid of stainless spoon, washed with soap and rinsed with distilled water for each sampling [12]. These roads are Adebayo road, Basin road, Ilawe road, Ajilosun road, Mathew road, Okeyinmi road, New Iyin road, Odo-Ado road, State secretariat road, University road. Two sampling sites were designated on each road. The samples were collected once every month for five months during the rainy season from May to September 2012. All the samplings were performed three days after the rain. Samples collected were stored in sealed polythene bags and transported to the laboratory for pre-treatment and analyses. Soil samples were air dried, mechanically ground using a stainless shell roller and served to obtain < 2 mm fraction. A 20-30 g sub sample was drawn from the bulk soil (< 2 mm fraction) and reground to obtain < 200 µm fraction using a mortar and pestle. The fire material used to determine the pH using soil water ratio of 1:5 using a Consort C862 bench top conductivity /pH/DO meter. Organic carbon was determined by the method Walkley and Black method. XRF analyses were carried out at the laboratory using hand held Thermo Scientific XLT-793 NITON energy-dispersive XRF analyzer having as excitation source a miniaturized 30 kV X-ray tube. Each soil sample was analyzed five times for 240 s using two X-ray filters, one for elements from K to Cu and the second for elements from Zn to Sb.

3. RESULT AND DISCUSSION

Soil pH values are presented in Table 1 indicating that the soils collected around Adebayo, New Iyin road, Basiri road, Mathew road, Ajilosun, road, Ilawe road, Okeyinmi road, Odo-Ado road, state secretariat road, and University road are alkaline (pH in the range of 7.973 ±0.05 to 8.846 ±0.12) and the control soil of Ado- Ekiti city is slightly acidic (pH = 6.185 ±0.05).

Table 1. Mean values pH for the investigated soil sample.

Soil sample	1	2	3	4	5	6	7	8	9	10	control
pH	8.846 ±0.12	8.837 ±0.07	8.662 ±0.14	8.409 ±0.03	8.783 ±0.05	8.484 ±0.02	8.442 ±0.06	8.557 ±0.10	7.973 ±0.05	8.373 ±0.22	6.185 ±0.07

XRF results for the collected soil samples evidenced the existence of the following major and minor elements: Fe, K, Mn, Ti (major) and Cr, Cu, Ni, and Zr (minor). The average concentrations of heavy metal Mn, Cr, Cu, Ni, and Zr of five measurements of each of the soil samples are given in Table 2. For the elements Ag, Cd, Hg, Sb, Se, and Sn the XRF results have not been reported because their concentrations were below the detection limits.

Table 2. Mean values of heavy metal content in the investigated soils sample.

Element	1	2	3	4	5	6	7	8	9	10	control
Fe	476.88	461.72	460.66	412.30	456.38	427.20	443.15	430.50	411.20	410.13	187.52
K	58.40	55.33	42.70	47.82	58.90	52.40	52.14	45.13	48.23	40.12	25.15
Mn	75.81	71.22	68.14	70.84	73.71	58.42	67.33	68.52	57.72	51.33	23.10
Ti	1.64	1.57	1.44	1.23	1.54	1.48	1.50	1.49	1.22	1.10	0.07
Cr	10.13	10.01	9.92	9.02	8.09	8.99	9.51	8.54	7.74	7.41	6.93
Cu	32.79	29.89	25.13	31.38	42.79	25.13	30.54	32.02	23.64	21.74	<15
Ni	62.63	65.55	60.88	58.7	61.41	62.67	56.46	53.30	47.83	47.18	41.93
Zr	1.85	1.23	0.62	0.85	1.14	0.91	0.41	0.23	0.31	0.35	0.09

1 - Adebayo road, 2 - Oke Iyinmi road, 3 - Ajilosun road, 4 - Mathew road, 5 - University road, 6 - Odo-Ado road, 7 - Ilawe road, 8 - Basiri road, 9 - State secretariat road, 10 - New Iyin road.

Total Fe concentrations in metal silt sediment ranged from 410.13 mg/kg at New Iyin road to 476.88 mg/kg at Adebayo road. Total Ti ranged from 1.101 mg/kg at New Iyin road to 1.728 mg/kg at Mathew road, ranged from 21.74 mg/kg to 42.79 mg/kg at New Iyin road and University road. Respectively ranged from 0.23 mg/kg to 1.85 mg/kg at Basiri roads and Adebayo road respectively. Ni ranged from 47.118 mg/kg at New Iyin road to 65.55 mg/kg at Oke Iyinmi road. Mn ranged from 51.33 mg/kg to 75.81 mg/kg at New Iyin road and Adebayo road as show in Table 2. The value of the metal at the road was higher than the control.

The three road, Adebayo Oke Iyinmi, and University road ranked highest in traffic density had the highest Ti, Cr, Ni, Zr and Mn Contents in soil, which were above the recommended mean for agriculture soil but lower than the maximum tolerable level proposed for agriculture soil (90-300 mg/kg), [13].

The mean and medium were used as estimates of central tendency standard error of the mean were all small. The distribution of original data for Fe, K, Mn, Ti, Cr, Cu, Ni and Zr are positive skewed. The substantial different in the symmetric parameter in the case of K, Ni, Cu, Fe, Mn and Ti indicate a non- normal distribution. This supporting a possibility of random infiltration of the metals from some anthropogenic source s. Large standard deviations in the case of Fe, Mn, Cu and N levels revealed their random fluctuating concentration level in the sediment. Among significant variable that controls or influences the distribution and concentration of heavy metal in the environment are the size of sediments and organic matter, Table 3 [14-16].

Table 3. Basic statistical parameters for the distribution of selected metals mg/kg in road metal silt sediment samples from Ado-Ekiti.

Element	Min	Max	Mean	Median	SD	SE
Fe	410.13	476.88	441.02	437.18	43.21	0.964
K	40.12	58.90	50.14	49.98	29.40	0.588
Mn	51.33	75.81	66.30	68.33	43.83	0.877
Ti	1.101	1.617	1.43	1.468	0.62	0.012
Cr	7.41	10.12	8.61	8.96	10.20	0.204
Cu	21.74	42.79	29.51	30.22	24.14	0.483
Ni	41.18	65.55	57.66	59.79	31.46	0.629
Zr	0.31	1.85	0.79	0.77	0.48	0.009

The degree of correlation between trace metal and organic matter and size distribution is often used to study the origin of many metals [17-27]. To verify this relationship in this study, correlations between all the metals and the parameter mention were carried out

4. CONCLUSION

XRF technique has been employed in order to establish the type of metal in soil sediment in Ado-Ekiti road. The experimental result indicate that the concentrations of heavy element varies from zone depend on the level of traffic volume on the road and population and they are greater than the level detected in the control soil. Anthropogenic release give rise to higher concentration of the metal relative to the normal background value and in some location their level exceed then alert level admitted by the Nigerian guideline.

References

- [1] Stone M., Marsalek J., *Air and Soil Pollution* 87 (1996) 149-169.
- [2] Pereira E., J. A. Baptista-Neto, B. J. Smith, J. J. Mcallister, *Ann. Braz. Acad. Sci.* 79 (2007) 739-750.
- [3] Sutherland R. A., C. A. Tolosa, *Environ. Pollut.* 110 (2000) 483-495.
- [4] Pagotto C., N. Remy, M. Legret, P. Le Cloirec, *Environ. Technol.* 22 (2001) 307-319..
- [5] Ferguson J. E., N. D. Kim, *Sci. Total Environ.* 100 (1991) 125-150.
- [6] Watts S. E. J., B. J. Smith, *Sci. Total Environ.* 146 (1994) 507-514.
- [7] McAllister J. J., B. J. Smith, J. A. Baptista Neto, *Environ. Geochem. Health* 22 (2000) 195-210.

-
- [8] McAllister J. J., B. J. Smith J. A. Baptista Neto, J. K. Simpson, *Environ. Geochem. Health* 27 (2009) 429-441.
- [9] McAllister J. J., B. J. Smith, J. A. Baptista Neto, J. K. Simpson, *Environ. Geochem. Health* 27 (2005) 429-441.
- [10] Pereira E., J. A. Baptista-Neto, B. J. Smith, J. J. Mcallister, *Ann. Braz. Acad. Sci.* 79 (2007) 739-750.
- [11] Sutherland R. A., *Environ. Pollut.* 121 (2003) 229-237.
- [12] Awofolu O. R., *Environ. Monitor. Assess.* 105 (2005) 431-447.
- [13] ICRCCL. (1987). *Interdepartmental committee for the Redevelopment of Contaminated Land, Guidance on the Assessment and Redevelopment of Contaminated Land*. Paper 59/83 2nd Edn. Department of the Environment, London
- [14] Lin Y. P., T. P. Teng, T. K. Chang, *Landscape Urban Plann.* 62 (2002) 19-35.
- [15] Huang K. M., S. Lin, *Chemosphere* 53 (2003) 1113-1121.
- [16] Lakhan V. C., K. Cabana, P. D. LaValle, *J. Coast. Res.* 19 (2003) 600-608.
- [17] Jumbe A. S., N. Nandini, *Am. J. Environ. Sci.* 5 (2009) 678-687.
- [18] Ebad Bashiri, Jahanbakhsh Bashiri, Farhad Karimi, *International Letters of Natural Sciences* 3 (2013) 7-20.
- [19] Daniszewski P., *International Letters of Chemistry, Physics and Astronomy* 3 (2012) 86-92.
- [20] Daniszewski P., *International Letters of Chemistry, Physics and Astronomy* 4 (2012) 112-118.
- [21] Daniszewski P., *International Letters of Chemistry, Physics and Astronomy* 5 (2012) 80-87.
- [21] Daniszewski P., Konieczny R., *International Letters of Chemistry, Physics and Astronomy* 4 (2013) 91-97.
- [22] Daniszewski P., Konieczny R., *International Letters of Chemistry, Physics and Astronomy* 8(3) (2013) 269-278.
- [23] Daniszewski P., Konieczny R., *International Letters of Chemistry, Physics and Astronomy* 8(3) (2013) 279-287.
- [24] Singare P. U., Talpade M. S., Dagli D. S., Bhawe V. G., *International Letters of Chemistry, Physics and Astronomy* 8(2) (2013) 94-104.
- [25] Piotr Daniszewski, *International Letters of Chemistry, Physics and Astronomy* 10(2) (2013) 218-226
- [26] Piotr Daniszewski, *International Letters of Chemistry, Physics and Astronomy* 12 (2013) 72-81
- [27] Piotr Daniszewski, *International Letters of Chemistry, Physics and Astronomy* 13 (2013) 13-22