

The second-connectivity and second-sum-connectivity indices of Armchair Polyhex Nanotubes $TUAC_6[m,n]$

Mohammad Reza Farahani

Department of Mathematics, Iran University of Science and Technology (IUST),
Narmak, Tehran 16844, Iran

E-mail address: Mr_Farahani@Mathdep.iust.ac.ir , mrfarahani88@gmail.com

ABSTRACT

The m -connectivity and m -sum connectivity indices of G are defined as to be ${}^m\chi(G) = \sum_{v_1v_2 \dots v_{m+1}} \frac{1}{\sqrt{d_{i_1}d_{i_2} \dots d_{i_{m+1}}}}$ and ${}^mX(G) = \sum_{v_1v_2 \dots v_{m+1}} \frac{1}{\sqrt{d_{i_1} + d_{i_2} + \dots + d_{i_{m+1}}}}$, where $v_1v_2 \dots v_{m+1}$ runs over all paths of length m in G and d_i is the degree of vertex v_i . In this paper, we give explicit formulas for the second-connectivity and second-sum-connectivity indices of an infinite class of Armchair Polyhex Nanotubes $TUAC_6[m,n]$.

Keywords: Molecular Graph; Armchair Polyhex Nanotubes; m -connectivity index; m -sum connectivity index

1. INTRODUCTION

Let $G = (V, E)$ be a simple connected graph with the vertex set $V(G)$ and the edge set $E(G)$. $|V(G)| = n$, $|E(G)| = e$ are the number of vertices and edges. The m -connectivity index of G is defined as

$${}^m\chi(G) = \sum_{v_1v_2 \dots v_{m+1}} \frac{1}{\sqrt{d_{i_1}d_{i_2} \dots d_{i_{m+1}}}},$$

where $v_1v_2 \dots v_{m+1}$ runs over all paths of length m in G and d_i is the degree of vertex v_i . In particular, 1-connectivity and 2-connectivity indices are defined as

$$\chi(G) = \sum_{e=uv \in E(G)} \frac{1}{\sqrt{d_u d_v}}$$

and

$${}^2\chi(G) = \sum_{v_{i_1}v_{i_2}v_{i_3}} \frac{1}{\sqrt{d_{i_1}d_{i_2}d_{i_3}}}$$

The 1-connectivity index (now called *Randić index*) introduced by *Milan Randić* in 1975 [1], where uv ranging over all pairs of adjacent vertices of G .

A closely related variant of the Randić connectivity index called the *sum-connectivity index* was introduced by *B. Zhou* and *N. Trinajstić* [2-7] in 2008 as the sum over all edges of the graph of the terms $(d_u + d_v)^{-1/2}$,

$$X(G) = \sum_{v_uv_v} \frac{1}{\sqrt{d_u + d_v}}$$

where d_u and d_v are the degrees of the vertices u and v , respectively.

Also, the m -sum connectivity index of G is defined as

$${}^mX(G) = \sum_{v_{i_1}v_{i_2}\dots v_{i_{m+1}}} \frac{1}{\sqrt{d_{i_1} + d_{i_2} + \dots + d_{i_{m+1}}}}$$

where $v_{i_1}v_{i_2}\dots v_{i_{m+1}}$ runs over all paths of length m in G .

In particular, *2-sum-connectivity index* is equal to

$${}^2X(G) = \sum_{v_{i_1}v_{i_2}v_{i_3}} \frac{1}{\sqrt{d_{i_1} + d_{i_2} + d_{i_3}}}$$

For more study about the m -connectivity index, the readers may consult [8-20] and references cited therein and some mathematical properties of the (general) sum-connectivity were given in [19-26].

In this paper, we give explicit formulas for the second-connectivity and second-sum-connectivity indices of an infinite class of Armchair Polyhex Nanotubes $TUAC_6[m,n]$.

2. RESULTS AND DISCUSSION

Consider the molecular graph Armchair Polyhex Nanotubes $TUAC_6$ and let we denote the number of hexagons in the first row/column of the 2D-lattice of $TUAC_6[m,n]$ (Figure 1) by m and n , respectively. For other related research and historical details, see the paper series [27-35] and the general representation of this nano structure is shown in Figure 1 and Figure 2.




Figure 1. The 2D Lattice (left) and 3D Lattice (right) of Armchair polyhex nanotubes of $TUAC_6[m,n]$.

The vertices/atoms and edges/bonds sets of the Armchair Polyhex Nanotubes $TUAC_6[m,n]$ are equal to $|V(TUAC_6[m,n])| = 2m(n+1)$ and $|E(TUAC_6[m,n])| = 3mn+2m$, respectively. Since $|V_2| = |\{v \in V(G) \mid d_v = 2\}| = 2m+2m$ and $|V_3| = |\{v \in V(G) \mid d_v = 3\}| = 2mn$.

Let we define d_{ijk} as a number of 2-edges paths with 3 vertices of degree i, j and k , respectively. It is obvious, $d_{ijk} = d_{kji}$ and an edge $e = v_i v_j$ is equal to d_{didj} .

Now, from Figure 2, we see that for every vertex in V_2 , there is a 2-edges path d_{223} (The red path in Figure 2) and also there are two 2-edges paths d_{233} (The yellow path in Figure 2).

From Figure 1, (the 2-dimensional lattice of Armchair Polyhex Nanotubes $TUAC_6[m,n]$, $\forall m, n > 1$), it's easy to see that the number of 2-edges paths in $TUAC_6[m,n]$ is equal to $6mn+4m$. Since there are six 2-edges paths for all hexagon or cycle C_6 in $TUAC_6[m,n]$ and also there is an addition 2-edges path for every member of V_2 . In other words,

$$d^2(TUAC_6[m,n]) = \underbrace{6 \times mn}_{\text{cycles } C_6} + \underbrace{4m}_{V_2}$$

So, from Figure 2, one can see that for all other vertices in V_3 , there are $d^2(TUAC_6[m,n]) - 4m - 2 \times 4m = 6mn - 8m$ 2-edges paths d_{333} (We marked these 2-edges paths by black color in Figure 2).




Figure 2. Examples of 2-edges paths d_{223} , d_{233} and d_{333} of $TUAC_6[m,n]$.

Now, by using above mentions, we have the second-connectivity index of *Armchair Polyhex Nanotubes* $TUAC_6[m,n]$ as

$$\begin{aligned}
 {}^2\chi(TUAC_6[m,n]) &= \sum_{v_i v_2 v_3} \frac{1}{\sqrt{d_{i_1} \times d_{i_2} \times d_{i_3}}} \\
 &= \frac{d_{223}}{\sqrt{2 \times 2 \times 3}} + \frac{d_{233}}{\sqrt{2 \times 3 \times 3}} + \frac{d_{333}}{\sqrt{3 \times 3 \times 3}} \\
 &= \frac{4m}{2\sqrt{3}} + \frac{8m}{3\sqrt{2}} + \frac{6mn - 8m}{3\sqrt{3}} \\
 &= \left(\frac{6n\sqrt{3} - 2\sqrt{3} + 12\sqrt{2}}{9} \right) m
 \end{aligned}$$

and also the 2-sum-connectivity index of $TUAC_6[m,n]$, $\forall m, n > 1$ is equal to

$$\begin{aligned}
 {}^2X(TUAC_6[m,n]) &= \sum_{v_i v_2 v_3} \frac{1}{\sqrt{d_{i_1} + d_{i_2} + d_{i_3}}} \\
 &= \frac{d_{223}}{\sqrt{2 + 2 + 3}} + \frac{d_{233}}{\sqrt{2 + 3 + 3}} + \frac{d_{333}}{\sqrt{3 + 3 + 3}} \\
 &= \frac{4m\sqrt{7}}{7} + \frac{4m\sqrt{2}}{2} + \frac{6mn - 8m}{3}
 \end{aligned}$$

$$= \left(2n + \frac{24\sqrt{7} + 84m\sqrt{2} - 112}{42} \right) m$$

Finally, from reference [16], we know that the first-connectivity index (Randić index) and first-sum-connectivity index of Armchair Polyhex Nanotubes $TUAC_6[m,n]$ are equal to

$$\chi(TUAC_6[m,n]) = \left(n + \frac{2\sqrt{6} + 1}{6} \right) m$$

and

$$X(TUAC_6[m,n]) = \left(\frac{\sqrt{6}n + 1}{2} + \frac{2\sqrt{5}}{5} - \frac{\sqrt{6}}{6} \right) m$$

3. CONCLUSION

In this report, we study some properties of two connectivity indices of (molecular) graphs that called second-connectivity and second-sum-connectivity indices and were defined as

$${}^2\chi(G) = \sum_{v_i v_i v_i} \frac{1}{\sqrt{d_i d_i d_i}}$$

and

$${}^2X(G) = \sum_{v_i v_i v_i} \frac{1}{\sqrt{d_i + d_i + d_i}}$$

where $\mathcal{V}_{i_1} \mathcal{V}_{i_2} \dots \mathcal{V}_{i_{m+1}}$ runs over all paths of length m in G and d_i is the degree of vertex v_i .

In continue, closed analytical formulas for these connectivity indices of a physico chemical structure of Hexagonal nanotubes are given. This nano structure is Armchair Polyhex Nanotubes $TUAC_6[m,n]$. The structures of Armchair Polyhex Nanotubes $TUAC_6[m,n]$ consist of several Hexagon C_6 .

References

- [1] M. Randić, *J. Am. Chem. Soc.* 97(23) 6609 (1975).
- [2] B. Zhou, N. Trinajstić, *J. Math. Chem.* 47 (2010) 210-218.
- [3] M. Randić, P. Hansen, *J. Chem. Inf. Comput. Sci.* 28 (1988) 60.
- [4] A. R. Ashrafi, P. Nikzad, *Digest. J. Nanomater. Bios.* 4(2) (2009) 269-273.
- [5] E. Estrada, *J. Chem. Inf. Comput. Sci.* 35 (1995) 1022.
- [6] E. Estrada, *Chem. Phys. Lett.* 312 (1995) 556.
- [7] B. Zhou, N. Trinajstić, *J. Math. Chem.* 46 (2009) 1252-1270.
- [8] Z. Mihali, N. Trinajstić, *J. Chem. Educ.* 69(9) (1992) 701.
- [9] D. Morales, O. Araujo, *J. Math. Chem.* 13 (1993) 95.
- [10] M. V. Diudea, P. E. John, *MATCH Commun. Math. Comput. Chem.* 44 (2001) 103.
- [11] M. V. Diudea, A. Graovac, *MATCH Commun. Math. Comput. Chem.* 44 (2001) 93.
- [12] M. V. Diudea, *MATCH Commun. Math. Comput. Chem.* 45 (2002) 109.
- [13] S. Yousefi, A. R. Ashrafi, *MATCH Commun. Math. Comput. Chem.* 56 (2006) 169.
- [14] X. Li, I. Gutman, *Mathematical Chemistry Monographs* 1 (2006) 330.
- [15] A. R. Ashrafi, P. Nikzad, *Digest Journal of Nanomaterials and Biostructures* 4(2) (2009) 269.
- [16] M. R. Farahani, *Acta Chim. Slov.* 59 (2012) 779-783.
- [17] M. R. Farahani, *Acta Chim. Slov.* 60 (2013) 198-202.
- [18] M. R. Farahani, M. P. Vlad, Some Connectivity Indices of Capra-Designed Planar Benzenoid Series $C_n(C_6)$. *Studia Universitatis Babeş-Bolyai Chemia*. (2014), In press.
- [19] M. R. Farahani, *Polymers Research Journal* 7(3) (2013).
- [20] M. R. Farahani, K. Kato, M. P. Vlad. *Studia Universitatis Babeş-Bolyai Chemia* 58(2) (2013) 127-132.
- [21] F. Ma, H. Deng, On the sum-connectivity index of cacti. *Mathematical and Computer Modelling*. February 2011.
- [22] B. Zhou, N. Trinajstić, *J. Math. Chem.* 47 (2010) 210-218.
- [23] Z. Du, B. Zhou, N. Trinajstić, *J. Math. Chem.* 47 (2010) 842-855.
- [24] Z. Du, B. Zhou, *On sum-connectivity index of bicyclic graphs*. arXiv:0909.4577v1.
- [25] Z. Du, B. Zhou, N. Trinajstić, *Appl. Math. Lett.* 2010, 24 (2010) 402-405.
- [26] R. Xing, B. Zhou, N. Trinajstić, *J. Math. Chem.* 48 (2001) 583-591.
- [27] M. V. Diudea, *MATCH, Commun. Math. Comput. Chem.* 45 (2002) 109-122.
- [28] A. R. Ashrafi, G. R. Vakili-Nezhaad, *Journal of Physics: Conference Series* 29 (2006) 181-184.
- [29] S. Yousefi, H. Yousefi-Azari, A. R. Ashrafi, M. H. Khalifeh, *JSUT* 33(3) (2008) 7-11.

- [30] A. Iranmanesh, Y. Alizadeh, *Digest. J. Nanomater. Bios.* 4 (2004) 607-611.
- [31] H. Shabani, A. R. Ashrafi, *Digest. J. Nanomater. Bios.* 4 (2009) 423-428.
- [32] S. Alikhani, M. A. Iranmanesh, *Digest. J. Nanomater. Bios.* 5 (2010) 1-7.
- [33] M. R. Farahani, *Journal of Advances in Physics* 3(1) (2013) 191-196.
- [34] M. R. Farahani, *Computing GA₅ Index of Armchair Polyhex Nanotube*, *Le Matematiche*, (2013), In press.
- [35] M. R. Farahani, *Proc. Rom. Acad., Series B*, 15(1) (2013) 3-6.

(Received 01 March 2014; accepted 07 March 2014)