Connective Eccentric Index of Circumcoronene Homologous Series of Benzenoid H_k

Mohammad Reza Farahani

Department of Applied Mathematics of Iran University of Science and Technology (IUST),
 Narmak, Tehran 16844, Iran

E-mail address: Mr_Farahani@Mathdep.iust.ac.ir , MrFarahani88@Gmail.com

ABSTRACT

Let G be a molecular graph, a topological index is a numeric quantity related to G which is invariant under graph automorphisms. The eccentric connectivity index $\xi(G)$ is defined as $\xi(G) = \sum_{v \in V(G)} d_v \times \varepsilon(v)$ where d_v , $\varepsilon(v)$ denote the degree of vertex v in G and the largest distance between v and any other vertex u of G . The connective eccentric index of graph G is defined as $C^\xi(G) = \sum_{v \in V(G)} \frac{d_v}{\varepsilon(v)}$. In the present paper we compute the connective eccentric index of *Circumcoronene Homologous Series of Benzenoid* H_k ($k \geq 1$).

Keywords: Molecular graphs; Benzenoid; Connective eccentric index; Eccentric connectivity index

1. INTRODUCTION

In theoretical chemistry molecular structure descriptor or topological indices, are used to compute properties of chemical compounds. Throughout this paper, graph means simple connected graph [1-3]. Let G be a molecular graph, the vertex and edge sets of a graph G are denoted by $V(G)$ and $E(G)$, respectively. If $x, y \in V(G)$ then the distance $d(x, y)$ between x and y is defined as the length of a minimum path connecting x and y .

In 1997, the *Eccentric Connectivity index* $\xi(G)$ of the molecular graph G was proposed by Sharma, Goswami and Madan and is defined as [4]:

$$\xi(G) = \sum_{v \in V(G)} d_v \times \varepsilon(v)$$

where d_v denotes the degree of the vertex v in G and $\varepsilon(v)$ denote the largest distance between v and any other vertex u of G . In other words, $\varepsilon(v) = \max\{d(v, u) \mid \forall v \in V(G)\}$.

In 2000, the *Connective Eccentric index* $C^\xi(G)$ was defined by Gupta, Singh and Madan [5,6] as follows:

$$C^\xi(G) = \sum_{v \in V(G)} \frac{d_v}{\varepsilon(v)}$$

where d_v , $\varepsilon(v)$ denote the degree and eccentric of vertex v in G . See [7-26] for more details and other versions of *Eccentric indices* and *Eccentric polynomials*.

The goal in this paper is computing the Connective eccentric index of *Circumcoronene Homologous Series of Benzenoid* H_k ($k \geq 1$).

2. RESULTS AND DISCUSSION

In this section, we compute the Connective eccentric index $C^\zeta(G)$ of Circumcoronene Homologous Series of Benzenoid. Three first members of this Benzenoid family ($H_1 = \text{benzene}$, $H_2 = \text{coronene}$ and $H_3 = \text{circumcoronene}$) are shown in Figure 1. Circumcoronene Homologous Series of Benzenoid is generated from famous molecule *Benzene* or cycle C_6 . We encourage reader to references [18-38] to study some properties of this Benzenoid family.

Figure 1. Three first members of Circumcoronene Homologous Series of Benzenoid: $H_1 = \text{benzene}$, $H_2 = \text{coronene}$ and $H_3 = \text{circumcoronene}$ [18-26].

$\forall k \in \mathbb{N}$ Circumcoronene Homologous Series of Benzenoid H_k has $6k^2$ atoms/vertices and $9k^2 - 6k$ bonds/edges (see Figure 2). For further study and more detail of this Benzenoid family, see the paper series [27-35]. Now, we have following theorem for this benzenoid graphs.

Theorem 1. Let G be the Circumcoronene Homologous Series of Benzenoid H_k ($\forall k \geq 1$). Then the Connective Eccentric index $C^\zeta(G)$ of H_k is equal to

$$C^\zeta(H_k) = \sum_{i=1}^{k-1} \left(\frac{9i(4k+4i-1)}{2i^2 + (4k-1)i + 2k^2 - k} \right) + \frac{12k}{4k-1}$$

Before prove the Theorem 1, we denote all vertices and edges of Circumcoronene Homologous Series of Benzenoid H_k , as follow and is shown in Figure 2, (Z_i , is the cycle finite group):

$$V(H_k) = \{ \gamma_{z,j}^i, \beta_{z,l}^i \mid i = 1, \dots, k, j \in Z_i, l \in Z_{i-1}, z \in Z_6 \}$$

$$\text{and } E(H_k) = \{ \beta_{z,j}^i \gamma_{z,j}^i, \beta_{z,j}^i \gamma_{z,j+1}^i, \beta_{z,j}^i \gamma_{z,j+1,1}^{i-1} \mid i \in \mathbb{Z}_k, j \in \mathbb{Z}_i, z \in \mathbb{Z}_6 \}$$

Proof. By considering Circumcoronene Homologous Series of Benzenoid $G = H_k$ ($\forall k \geq 1$) as shown in Figure 2 and refer to [18-26] and using the *Ring-cut Method* for circumcoronene homologous series of Benzenoid, we can compute its connective eccentric index. The *Ring-cut Method* is a modify version of the thoroughbred *Cut Method*. The general form of this method is introduced in [18-26] For more study and detail information of the Cut Method see [28,31,32].

Figure 2. The general representation of Circumcoronene Homologous Series of Benzenoid H_k ($k \geq 1$) [18-26].

To compute the connective eccentric index of H_k , we see that

$$\forall i = 2, \dots, k; j \in \mathbb{Z}_{i-1} \text{ \& } z \in \mathbb{Z}_6: \varepsilon(\beta_{z,j}^i) = 2k + 2i - 2$$

$$\forall i = 1, \dots, k; j \in \mathbb{Z}_i \text{ \& } z \in \mathbb{Z}_6: \varepsilon(\gamma_{z,j}^i) = 2k + 2i - 1$$

Also, by according to Figure 2, one can see that the vertices in general representation of molecular graph Circumcoronene Homologous Series of Benzenoid H_k have degree two or three, such that

$$V_2(H_k) = \{v \in V(H_k) | d_v = 2\} = \{\gamma_{z,i}^k | \forall i \in \mathbb{Z}_i \text{ \& } z \in \mathbb{Z}_6\}$$

and alternatively $V_3(H_k) = V(H_k) - V_2(H_k)$.

$$\begin{aligned} C^\xi(H_k) &= \sum_{v \in V(G)} \frac{d_v}{\varepsilon(v)} \\ &= \sum_{\gamma_{z,j}^k \in V_2(H_k)} \frac{d_{\gamma_{z,j}^k}}{\varepsilon(\gamma_{z,j}^k)} + \sum_{\gamma_{z,j}^i \in V_3(H_k)} \frac{d_{\gamma_{z,j}^i}}{\varepsilon(\gamma_{z,j}^i)} + \sum_{\beta_{z,j}^i \in V_3(H_k)} \frac{d_{\beta_{z,j}^i}}{\varepsilon(\beta_{z,j}^i)} \\ &= \sum_{\substack{\gamma_{z,j}^k \in V_2(H_k) \\ j \in \mathbb{Z}_6; z \in \mathbb{Z}_6}} \frac{2}{4k-1} + \sum_{\substack{\gamma_{z,j}^i \in V_3(H_k); z \in \mathbb{Z}_6 \\ i=1, \dots, k-1; j \in \mathbb{Z}_i}} \frac{3}{2k+2i-1} + \sum_{\substack{\beta_{z,j}^i \in V_3(H_k); z \in \mathbb{Z}_6 \\ i=2, \dots, k; j \in \mathbb{Z}_{i-1}}} \frac{3}{2k+2i-2} \\ &= \sum_{z=1}^6 \sum_{j=1}^k \left(\frac{2}{4k-1} \right) + \sum_{z=1}^6 \sum_{i=1}^{k-1} \sum_{j=1}^i \left(\frac{3}{2k+2i-1} \right) + \sum_{z=1}^6 \sum_{i=2}^k \sum_{j=1}^{i-1} \left(\frac{3}{2k+2i-2} \right) \\ &= 6k \left(\frac{2}{4k-1} \right) + \sum_{i=1}^{k-1} \left(\frac{3 \times 6i}{2k+2i-1} \right) + \sum_{i=2}^k \left(\frac{3 \times 6(i-1)}{2k+2i-2} \right) \\ &= \frac{2 \times 6k}{4k-1} + \sum_{i=1}^{k-1} \left(\frac{3 \times 6i}{2k+2i-1} \right) + \sum_{j=1}^{k-1} \left(\frac{3 \times 6j}{2k+2j} \right) \\ &= \sum_{i=1}^{k-1} \left(\frac{18i(2k+2i-1+2k+2i)}{2(k+i)(2k+2i-1)} \right) + \frac{12k}{4k-1} \end{aligned}$$

Thus $\forall k \geq 1$, the connective eccentric index of H_k is equal to

$$C^\xi(H_k) = \sum_{i=1}^{k-1} \left(\frac{9i(4k+4i-1)}{2i^2 + (4k-1)i + 2k^2 - k} \right) + \frac{12k}{4k-1}$$

and this completed the proof of Theorem 1.

3. CONCLUSION

The eccentric connectivity index $\xi(G)$ is defined as $\xi(G) = \sum_{v \in V(G)} d_v \times \varepsilon(v)$ where d_v , $\varepsilon(v)$ denote the degree of vertex v in G and the largest distance between v and any other vertex u of G . In this paper, we counting the connective eccentric index $C^\xi(G) = \sum_{v \in V(G)} \frac{d_v}{\varepsilon(v)}$ of *Circumcoronene Homologous Series of Benzenoid* H_k ($k \geq 1$).

References

- [1] H. Wiener, *J. Am. Chem. Soc.* 69(1) (1947) 17-20.
- [2] I. Gutman, N. Trinajstić, *Chem. Phys. Lett.* 17 (1972) 535-538.
- [3] R. Todeschini, V. Consonni, *Handbook of Molecular Descriptors*, Weinheim, Wiley-VCH, (2000).
- [4] V. Sharma, R. Goswami, A. K. Madan, *J. Chem. Inf. Comput. Sci.* 37 (1997) 273-282.
- [5] S. Gupta, M. Singh, A. K. Madan, *J. Mol. Graph. Model.* 18 (2000) 18-25.
- [6] S. Gupta, M. Singh, A. K. Madan, *J. Math. Anal. Appl.* 266 (2002) 259-268.
- [7] A.R. Ashrafi, M. Ghorbani, M. Hemmasi, *Digest. J. Nanomater. Bios.* 4(3) (2009) 483-486.
- [8] S. Alikhani, M. A. Iranmanesh, *Digest. J. Nanomater. Bios.* 6(1) (2011) 253-257.
- [9] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, New York, (1986).
- [10] M. A. Johnson, G. M. Maggiora, *Concepts and Applications of Molecular Similarity*, Wiley Interscience, New York, (1990).
- [11] M. Ghorbani, M. Ghazi, *Digest. J. Nanomater. Bios.* 5(4) (2010) 1107-1111.
- [12] V. Kumar, S. Sardana, A. K. Madan, *J. Mol. Model.* 10 (2004) 399-407.
- [13] M. Fischermann, A. Homann, D. Rautenbach, L. A. Szekely, L. Volkmann, *Discrete Appl. Math.* 122 (2002) 127-137.
- [14] S. Sardana, A. K. Madan, *MATCH Commun. Math. Comput. Chem.* 43 (2001) 85-89.
- [15] A. R. Ashrafi, M. Ghorbani, M. Jalali, *Optoelectron. Adv. Mater.-Rapid Comm.* 3 (2009) 823-826.
- [16] H. Dureja, A. K. Madan, *Med. Chem. Res.* 16 (2007) 331-341.
- [17] Z. Yar Ahmadi, *Iran. J. Math. Chem*, 1(2) (2010) 105-110.
- [18] M. R. Farahani. *Annals of West University of Timisoara-Mathematics and Computer Science* 51(2) (2013) 29-37.
- [19] M. R. Farahani. *Journal of Advances in Physics* 2(1) (2013) 48-52.
- [20] M. R. Farahani. *Int. J. Chem. Model.* 6(1) (2014).
- [21] M. R. Farahani, *Journal of Chemica Acta* 2 (2013) 22-25.

-
- [22] M. R. Farahani, *World Applied Sciences Journal* 21(9) (2013) 1260-1265.
- [23] M. R. Farahani, Two New Version of Zagreb Index of Circumcoronene Series of Benzenoid. *Int. J. Computational Sciences & Applications*. (2013) In press.
- [24] M. R. Farahani, *Chemical Physics Research Journal* 6(1) (2013) 27-33.
- [25] M. R. Farahani, *Journal of Chemica Acta* 2 (2013) 26-31.
- [26] M. R. Farahani, *Journal of Mathematical Nanoscience* 2(1) (2012) 15-20.
- [27] S. Klavžar, I. Gutman, B. Mohar, *J. Chem. Int Comput. Sci.* 35 (1995) 590-593.
- [28] V. Chepoi, S. Klavžar, Distances in benzenoid systems: Further developments. *Discrete Mathematics* 192 (1998) 27-39.
- [29] P. Zigert, S. Klavžar, I. Gutman, *ACH Models Chem.* 137 (2000) 83-94.
- [30] A. Soncini, E. Steiner, P. W. Fowler, R. W. A. Havenith, L. W. Jenneskens, *Chem. Eur. J.* 9 (2003) 2974-2981.
- [31] P. E. John, P. V. Khadikar, J. Singh, *J. Math. Chem.* 42(1) (2007) 27-45.
- [32] S. Klavžar, *MATCH Commun. Math. Comput. Chem.* 60 (2008) 255-274.
- [33] M. R. Farahani, M. P. Vlad, *Studia Univ. Babes-Bolyai, Chemia.* 57(4) (2012).
- [34] M. R. Farahani, *J. Applied Mathe. & Informatics.* 31(3) (2013).
- [35] M. R. Farahani, *Int. J. Chem. Model.* 6(1) (2014).
- [36] M. R. Farahani, *International Letters of Chemistry, Physics and Astronomy* 11(1) (2014) 74-80.
- [37] M. R. Farahani, *International Letters of Chemistry, Physics and Astronomy* 12 (2014) 56-62.
- [38] M. R. Farahani, *International Letters of Chemistry, Physics and Astronomy* 12 (2014) 63-68.

(Received 04 April 2014; accepted 12 April 2014)