Reusable Nanocatalyst: Zirconia and Sulfated Zirconia

Mitesh B. Gondaliya*, Tushar Maheta, Mayank J. Mamtora, Manish K. Shah

Department of Chemistry, Saurashtra University, Rajkot, Gujarat, India

*E-mail address: miteshgondaliya@scientist.com

ABSTRACT

1,5-benzodiazepines have been synthesized by Nano Zirconia and Nano Sulfated Zirconia in solvent free conditions. Synthesis was carried out by conventional as well as microwave method. Benzodiazepines were synthesized by reacting o-phenylenediamine and few ketones in the presence of catalytic amount of Nanoparticles. Particles used were sized in between 53-100 nm. Nanoparticles were characterized by FT-IR, DLS and powder XRD analysis.

Keywords: Nanocatalyst; Nanoparticles; Sulfated Zirconia; Synthesis; Benzodiazepines; Solid catalyst; Solvent free; Green Chemistry

1. INTRODUCTION

Benzodiazepines is very well known drug which gives various pharmacological importance, for example anxiolytic, anticonvulsant, hypnotic and sedative agent [1,2]. So benzodiazepines were of great interest in industries. There were many methods known, which were considered for up-gradation in urgencies. Many catalyst reported in literatures with significant boundaries were Silica supported 12-tungstophosphoric acid [3], boron trifluoride ether complex [4], Ytterbium triflate [5], Magnesium oxide/phosphorus oxychloride [6]. All methods in literature has its distinct use and also has some limitations like green approach, reusability, etc. The most prominent method is to use solid superacid catalyst, which can be further be recovered and reused several times. While, if we use nanomaterial, which have greater surface area compared to macro particles, may give faster reaction. So, taking in mind, we have used Nano Zirconia and Nano sulfated Zirconia for the synthesis of benzodiazepines.

2. EXPERIMENTAL

Thin-layer chromatography was accomplished on 0.2 mm precoated plates of silica gel G60 F254 (Merck). IR spectra were recorded on a FTIR-8400 spectrophotometer using DRS prob. ^1H (400 MHz) and ^{13}C (100 MHz) NMR spectra were recorded on a Bruker ADVANCE

¹H NMR spectrometer in DMSO-d₆. Chemical shifts are expressed in δ ppm downfield from TMS as an internal standard. Solvents were evaporated with a BUCHI rotary evaporator. Melting points were measured in open capillaries and are uncorrected. X-ray diffraction (XRD) measurements were performed on the XPERT-MPD by an X-ray diffractometer using Cu K α radiation source ($\lambda = 1.5418$) at power 200 kW. XRD Crystallite Size calculation was done using Scherrer equation from powder XRD data. Particle size measurements were carried out on DLS Nanotracer of Microtrac Europe GmbH. Domestic microwave was used at 400 W power. Preparation of Zirconia and Sulfated Zirconia were adapted^[7,8] and modified.

2. 1. Preparation of zirconia catalyst

Hydrated Zirconium oxychloride was dissolved in water then added ammonium hydroxide till solution became basic. The precipitate obtained was filtered. Zirconium-hydroxide gel was dried in oven then formed solid was crushed. The fine powder was calcinated at 600 °C for 4 hour. The particles obtained were stored in cold and anhydrous environment.

2. 2. Preparation of Sulfated-Zirconia

Previously prepared Zirconium hydroxide was acidified with 0.1 M sulfuric acid then stirred for 24 hour then centrifuged and dried in oven then formed solid was crushed. The fine powder was calcinated at 600 °C for 4 hour. The particles obtained were characterized by XRD, DLS and FT-IR.

2. 3. General procedure for synthesis of Benzodiazepines

A mixture of o-phenylenediamine (1 mmol), different ketone (2.5 mmol) and a catalytic amount of nanomaterial (0.02 mg) in round bottom flask were stirred at ambient condition. Reaction was monitored by TLC in intervals. After completion of reaction the nanoparticles were recovered by adding dichloromethane (10 ml) and then centrifuged in high RPM until nanoparticles were settled in bottom. Washed nanoparticles 2-3 times with dichloromethane and nanoparticles were reused in reaction wet/dried. Similarly reactions were carried out in microwave.

Scheme 1. General reaction for synthesis of 1,5-benzodiazepine derivatives.

2,2,4-trimethyl-2,3-dihydro-1H-benzo[*b*][1,4]diazepine (AZ-1): Yellow solid, mp 150-152 °C; Mass $M^+ = 188$; IR (KBr) $\nu(\text{cm}^{-1})$, 3330 -NH (s, br); ¹H NMR (δ ppm) 1.242 (s, 6H),

2.154 (s, 2H), 2.230 (s, 3H), 4.712 (s br, 1H), 6.825-6.965 (m, 4H); ^{13}C NMR (δ ppm) 170.76, 146.61, 139.33, 127.37, 126.82, 125.91, 121.08, 119.92, 79.29, 66.70, 45.15, 39.06, 29.38.

4,4'-(2-methyl-2,3-dihydro-1H-benzo[b][1,4]diazepine-2,4-diyl)diphenol (AZ-2): Yellow solid, mp 218-220 °C; Mass $M^+ = 344$; IR (KBr) $\nu(\text{cm}^{-1})$, 3330 -NH (s,br); ^1H NMR (δ ppm) 1.630 (s, 3H), 2.471 (d, 1H), 4.712 (brs, 1H), 8.4 (s,1H), 8.87 (s,1H), 6.3-6.7 (m, 4H), 6.8 -7.2 (m, 8H); ^{13}C NMR (δ ppm) 195.99, 170.30, 162.04, 155.71, 139.59, 228.79, 125.36, 115.17, 77.43, 59.76, 39.62, 30.18, 27.74, 20.61.

10-Spirocyclopentan-1,2,3,9,10,10a-hexahydrobenzo[b]-cyclopenta[e][1,4]diazepine (AZ-3): Yellow solid, mp 138-140 °C; Mass $M^+ = 268$; IR (KBr) $\nu(\text{cm}^{-1})$, 3330 -NH (s,br); ^1H NMR (δ ppm) 1.2-1.8 (m, 19H), 3.98 (s br, 1H), 6.89-7.25 (m, 4H); ^{13}C NMR (δ ppm) 178.27, 134.55, 145.08, 55.57, 126.59, 114.79, 121.80, 129.10, 43.54, 34.98, 37.07, 37.09, 22.56, 27.48, 22.51, 22.53, 25.1, 22.68.

2-methyl-2,4-diphenyl-2,3-dihydro-1H-benzo[b][1,4]diazepine (AZ-4): Yellow solid, mp 158-160 °C; Mass $M^+ = 312$; IR (KBr) $\nu(\text{cm}^{-1})$, 3330 -NH (s,br); ^1H NMR (δ ppm) 1.45 (s, 3H), 2.21 (s, 2H), 4.02 (s br, 1H), 6.89-7.37 (m 14H); ^{13}C NMR (δ ppm) 141.99, 149.90, 129.23, 125.24, 126.65, 114.88, 128.20, 126.10, 128.24, 121.9, 128.80, 128.44, 131.99, 125.80, 166.26, 80.81, 43.44, 29.40.

Table 1. Synthesis of 1,5-benzodiazepine derivatives.

Code	Reactant	Product	Yield % ^a	Time (In min)	
				Δ	MW
AZ-1	Acetone		95	40	2
AZ-2	p-Hydroxy acetophenone		85	50	4
AZ-3	Cyclohexanone		90	50	4

3. RESULT AND DISCUSSION

In this current work, we have synthesized Nano zirconium dioxide (ZrO_2) and sulfated zirconia. Both nanoparticles were studied for its catalytic activity to synthesis benzodiazepines derivatives. Nano particles were synthesized by solution gel method which is beneficiary in terms of bulk synthesis of Nano particles. The shape of Nano zirconium dioxide particles found from powder XRD were monoclinic and of Nano sulfated zirconia (SZ) particles were triclinic in shape. In particular, characterization we found that, in DLS, the size of nano zirconium dioxide was near about 100 nm, and the similar size of Nano sulfated zirconia was also found near 100 nm. From Powder XRD we found that the average bulk size of Nano zirconium dioxide was ranging between the 53-59 nm and geometrical structure was monoclinic. While, the average bulk size of Nano SZ was ranging between the 53-59 nm and geometrical structure was triclinic.

Summary

Data	Value	%Tile	Size(nm)
MV(nm):	366.0	95.00	341.0
MN(nm):	137.7	90.00	261.6
MA(nm):	278.9	80.00	139.4
CS:	21.51	60.00	111.8
SD:	39.90	50.00	105.6
PDI:	0.931	40.00	100.4
Mz:	363.9	30.00	95.80
σ :	155.0	20.00	91.30
Ski:	64.39261	10.00	86.10
Kg:	1,060	5.00	81.90

Figure 1. DLS of ZrO_2 .

Figure 2. DLS of Sulfated ZrO_2 .

Normal reaction time without any catalyst for the synthesis of Benzimidazole was around 24 hour, but when we add little amount of ZrO_2 powder reaction time was decreased to 90-100 minutes and yield was increased somewhat. While, activated Sulfated Zirconium dioxide powder were used in synthesis of benzimidazole, time of reaction was decrease to 40-60 minutes and yield was high in compared to other similar catalysts which were referenced in introduction chapter.

After completion of reaction, rest of solution were treated with dimethyl chloride to dissolve all precipitated organic part, then centrifuged the solution to recover the Nano particles by decantation of solution. Particles were washed twice by cold dimethyl chloride before its further use.

Table 2. Reusability of ZrO_2 and Sulfated ZrO_2 .

Sr. No.	Nanocatalyst	Average recovery ZrO_2	Average recovery Sulfated ZrO_2
1	1 st Run	80 %	85 %
2	2 nd Run	75 %	80 %
3	3 rd Run	70 %	76 %
4	4 th Run	65 %	72 %

Figure 3. Powder XRD of ZrO_2 and Sulfated ZrO_2 .

Figure 4. FT-IR of (A) ZrO₂ (B) Sulphated ZrO₂.

3. CONCLUSION

We were able to synthesize ZrO_2 and Sulfated ZrO_2 (SZ) Nanoparticles by Sol-Gel method. The nanoparticles were characterized by IR, Powder XRD and DLS. The sizes of Nanoparticles were found in the range of ~50-100 nm. The prepared Nanoparticles shows higher catalytic activity compared to other methods described. Synthesis of benzodiazepines derivatives were characterized by Mass, FT-IR and NMR. However, using this SZ Nanoparticles shows greater reusability up till 3-4 times, hence beneficiary as catalyst.

Acknowledgement

Authors are thankful for UGC-BSR fellowship. Also acknowledge Physics department for pXRD data and NFDD, Rajkot for NMR data.

References

- [1] D. J. Abraham, *Burger's Medicinal Chemistry and Drug Discovery*, John Wiley & Sons, Inc., Hoboken, NJ, USA, 2003.
- [2] R. G. Priest, U. V. Filho, R. Amrein, M. Skreta, Eds., *Benzodiazepines*, Springer Netherlands, Dordrecht, 1980.
- [3] M. Abdollahi-Alibeik, I. Mohammadpoor-Baltork, Z. Zaghaghi, B. H. Yousefi, *Catalysis Communications* 9 (2008) 2496.
- [4] J. A. L. Herbert, H. Suschitzky, *Journal of the Chemical Society, Perkin Transactions 1* (1974) 2657.
- [5] M. Curini, F. Epifano, M. C. Marcotullio, O. Rosati, *Tetrahedron Letters* 42 (2001) 3193.
- [6] M. . Balakrishna, B. Kaboudin, *Tetrahedron Letters* 42 (2001) 1127.
- [7] J. R. Sohn, H. W. Kim, *Journal of Molecular Catalysis* 52 (1989) 361.
- [8] X. SONG, A. SAYARI, *Catalysis Reviews* 38 (1996) 329.

(Received 02 July 2014; accepted 12 July 2014)