Nucleic Acid Content in Male Fresh Water Fish *N. notopterus* Exposed to Copper Sulphate

K. Ravikiran*, R. S. Kulkarni

Fish Endocrinology Research Laboratory
Department of Post Graduate Studies and Research in Zoology
Gulbarga University Gulbarga - 585 106 Karnataka, India

*E-mail address: ravikiranrattimani@gmail.com

ABSTRACT

The nucleic acid content in different tissues such as brain, liver, kidney & testis of both control and copper sulphate (CuSO_4) exposed freshwater fish *N. notopterus* has been studied. The following observation is made in both control and copper sulphate (CuSO_4) exposed fish. The testis contain large amount of DNA in comparison to other tissues. The degree of DNA content in control and copper sulphate (CuSO_4) exposed fish testis >liver>brain>kidney. The RNA content also exhibited similar to that of DNA, having higher amount in the testis. The degree of RNA content in control and copper sulphate (CuSO_4) exposed fish testis >liver>brain>kidney. The nucleic acid content of tissues get reduced under copper sulphate (CuSO_4) exposed in the male freshwater fish *N. notopterus* indicating copper sulphate as a pollutant effect the nucleic acid content in the tissue.

Keywords: DNA; RNA; *N. notopterus*; CuSO_4

1. INTROUDCTION

Water is an important natural resource essential for all living beings. The important water resources are ponds, lakes, rivers streams, etc. These water bodies provide water for drinking, irrigation, cooling industrial machines, and for other domestic purposes. So any harmful change in the water causes many ill-effects on the life of man and other animals. The undesirable change in water that has harmful affect on the life of man and domestic plants and animals is called water pollution. Urbanization due to population explosion and industrialization are the main causes for water pollution. Many organic and inorganic wastes from industries are discharged into water bodies. These wastes include heavy metals such as Hg, Cu, Zn lead etc., detergents, acids, alkalis, oils, phenols, petroleum, carbonates, alcohols, cyanides, arsenic, chlorine, etc.

Among several elements of the periodic table, there are 35 metals are associated with community and occupational exposure. Out of these, 23 are described as heavy metals. These elements are generally released in small amounts into the environment by processes like weathering of rocks, volcanic eruptions etc. and their intake/exposure is necessary in trace amounts for good health. But, presently, there is a steady increase in their concentration in all habitats owing to mining, electroplating, paints and dye, battery making industries etc. The release is rapid with the rapidly growing technology and heavy metal application in these

industries. The two most important factors that contribute to the deleterious effects of heavy metals as pollutants are their indestructible nature through bioremediation unlike organic pollutants and their tendency to accumulate in environment especially in the bottom sediments of aquatic habitats in association with organic and inorganic matter. Out of the several heavy metals in the industrial waste streams (Forstner and Prosi, 1979).

Aquatic environments are loaded with several types of organic and inorganic pollutants. Huge amounts of agriculture pesticides, used for crop protection, eventually enter into the aquatic system. Similarly, heavy metals, which are released as industrial effluents form the major constituents of aquatic pollution. The presence of excess quantities of these toxic pollutants in water bodies has caused mass mortality of fishes in the past (Wanganeo *et al.*, 1994).

The present study was undertaken on the freshwater fish *N. notopterus* keeping in view the above background. The study describes the toxicity levels of heavy metal like copper sulfate and their effects, of Copper sulphate found to be a pollutant causing deleterious effects on aquatic organisms at different levels, in the present study effects of copper sulphate on nucleic acid content of some important tissues of male freshwater fish *N. notopterus* has been undertaken.

2. MATERIALS AND METHODS

Fresh water fish *N. notopterus* (80-90 gm body weight) were brought from Bheema River around 40 kms away from Gulbarga. Fish were acclimatized for laboratory condition for 7 days before the beginning of the experiment as where fish were fed during this adaptation period. This supplementary aeration provided optimum dissolved oxygen. Dissolved oxygen levels were recorded before experiment and after experiment. Water temperature was measured before and after the completion of experiment. Stable temperature of 27 ± 3 is an optimum temperature during the whole period of study. Fish were fed with earthworm and boiled egg pieces once daily from the day of arrival until the end of the experiment. This light provided for all aquaria during adoption and was about 12 hrs of light: 12 hrs of dark/day during the days of study.

2.1. Use of Copper sulphate as a pollutant

2.1.1. Copper sulphate

It is classified as general used material by environmental protection agency. Copper sulphate is a fungicide used to control bacterial and fungal diseases of fruit vegetable, nuts and field crops. Some of the diseases that are controlled by this fungicide include mildew, leaf spots, blights and apple scab. It is also used as an algacide in irrigation and municipal water treatment system copper sulphate is an naturally occurring in-organic salt and copper is an essential trace element in plant and animal nutrition. It is available in dust wettable powders and fluid concentrates.

2.1.2. Physical properties

It is blue in colour and odorless, it gives off moisture when exposed to air, copper sulphate (CuSO_4) is highly corrosive to plain steel, iron and galvanized. It is indefinitely stable when kept dry and is stable to heat, cold or light burning copper sulphate may produce irritating or poisonous gases and pollution may be caused by runoff from fire control or dilution water.

2.1.3. Determination of LC_{50}

About 50 fishes from the stock were used for the determination of LC_{50} . The fish were divided into 5 groups containing 10 fish in each Group, Group-I was kept as control and group II, III, IV, V were exposed in different concentration of 15 ppm, 20 ppm, 25 ppm, 30 ppm separately in Aquaria containing 25 liter of water. For a period of 96 hrs duration the experimental solution were changed every 24 hrs. The concentration that kills 50% percent of the fish in 96hrs duration ($\text{LC}_{50}/96$ hrs) was determine by Static Bioassay method (Dandoroff *et al.*, 1951) by using the mortality values for 96 hrs the LC_{50} was determined by plotting the graph. The exposed fishes were sacrificed after 96 hrs (LC_{50}) and the tissues such as brain, liver, kidney and gonads were dissected out and processed for determination of the nucleic acid.

The nucleic acid (DNA and RNA) content of the tissue was estimated by Schneider (1940). In all the cases six observation were made and the results were expressed as arithmetic mean with their standard deviation, standard error and student "t" were made as described by Suedecur (1946) and Fisher (1963).

3. RESULTS AND DISCUSSION

After exposing fish, *N. notopterus* to different concentration of copper sulphate, the LC_{50} was found to be 25 ppm under this concentration (Table-1; Fig. 1) the fish were exposed for 96 hrs and sacrificed. The nucleic acid (DNA/RNA) content in different tissues such as brain, liver, kidney and testis of both control and copper sulphate (CuSO_4) exposed male freshwater fish *N. notopterus* has been studied, during September month (i.e., spawning phase of the reproductive cycle).

The observation is made in both control and copper sulphate (CuSO_4) exposed fish. Amongst the four tissues, the testis contains larger amount of DNA in comparison to other tissues. The degree of DNA content in control and copper sulphate (CuSO_4) exposed fish although remains same, the DNA content reduces after CuSO_4 exposure in all the tissues (testis > liver > brain > kidney). The RNA content also exhibited similar to that of DNA. The degree of RNA content in control and copper sulphate (CuSO_4) exposed fish although remains same, the RNA content reduces after CuSO_4 exposure (testis > liver > brain > kidney) in all the tissues.

The DNA/RNA ratio's in different tissues found to be in a control unexposed fish is (Brain 2:1, Liver 2:1, Testis 2:1 and only in Kidney it is 1:1), and in Copper sulphate (CuSO_4) exposed although remains same, it is in the reduced condition (Brain has 2:1, Liver 2:1, Testis 2:1, and only in Kidney it is 1:1). (Table-2, Fig-2).

The level of DNA and RNA was found to be decreased in brain, liver, gonads, and kidney of *N. notopterus* fish due to exposure to copper sulphate indicating toxicity effects

nucleic acid synthesis. The rapid histolysis in tissues because of copper sulphate treatment to fish is the only possible reason for this, alternation. Impairment of nucleic acid metabolism and the degradation of cells, resulting in the reduction of DNA content. Copper sulphate, inhibition of DNA changes, thus, might affect both protein as well as amino acid levels by decreasing the level of RNA in protein synthesis machinery. Copper sulphate as a potential inhibitor of DNA synthesis, might result in the reduction in the RNA level because of toxicity in nature many enzymes responsible for normal metabolic pathway are also affected.

RNA plays significant role in protein synthesis; depletion in RNA contents also results in depletion of protein levels. Hence, there is decrease in RNA level thus reducing protein synthesis. This is because of the animals requires more energy to overcome the stress upon exposure to the pollutant.

The DNA and RNA content of male fish tissues gets reduced under copper sulphate (CuSO_4) exposure in the freshwater fish *N. notopterus* indicating copper sulphate (CuSO_4) as a pollutant effects the reduction in the nucleic acid content in the tissues.

Environmental problems have always existed throughout human history but widespread recognition has come, understandably, one belatedly after many year of steadily accumulating pollution, pollution of environment by heavy metals is of prime importance unrestrained release of heavy metals into environment via discharge of industrial effluent sewage and agro-chemicals into the water source has not only rendered in unusable but at the same time has produced great harm to fish (Vineetashukla *et al.*, 2002).

The nucleic acid (DNA/RNA) content in different tissue of *N. notopterus* exposed to copper sulphate (CuSO_4) at LC_{50} concentration for 96 hrs indicate that the DNA/RNA content of tissues reduced. Indicating copper sulphate as a pollutant effects the nucleic acid content in the tissues.

Adult pre-spawning fish *Labeo rohita* were sub lethally ($1/5^{\text{th}}$ 96 hrs LC_{50}) exposed to mercuric chloride and metacid-50 (methyl parathion). Accumulation of mercury and methyl parathion was studied and it was found that pre-spawning ovary appears as a potent organ for deposition of both the pollutants. RNA/DNA ratio of control and treated fish were studied. It was found that the significant decrease in RNA/DNA ratio occurs after 9 and 30 days of exposure for mercury and 30 days for methyl parathion. Fluorescence microscope studies by alcidine orange staining method were also performed to show how much is it related to biochemical alterations. In some cases loss of metachromasia is correlated with the fall in RNA/DNA ratio. Some other abnormalities like fall in stage II and I oocyte ratio and necrosis was also observed (Ajitkumar *et al.*, 2002).

Accumulation of heavy metals and pesticide residues and their effect on fish physiology have been reported by several workers (Wilder and Stanley, 1989). Bhattacharya and Pandey (1989) reported Sonatox induced changes in the ovaries of *Oreochromis mossambicus*. Rajnarayan and Sathyanesan (1986) reported mercurial induced changes in the ovarian development of *Channa punctatus*. Sahai (1989) reported some histopathological changes in the ovaries of *Puntius ticto* by the treatment of several insecticides.

RNA/DNA ratio indicates the degree of metabolic (protein) synthesis. It is based on the fact that the DNA content per cell is constant within the same species, and the RNA is mainly ribosomic and varies with the rate of protein synthesis. It has been traditionally used as a growth index in ecological studies, in aquaculture and as a biomarker in long-term exposure experiments in ecotoxicology (Miliou *et al.*, 1998). The sensitivity of the ratio as a biomarker has been questioned based on its lack of response in some experiments (Benton *et al.*, 1994; Kim and Kang, 2004). In a recent study, Vidal *et al.* (2006) were able to associate RNA/DNA variations in 24 hrs period and at different hours of the day in starved para larvae of the

loliginid squid. The larval stages of cephalopods are very fast growing forms, metabolically very active. The sensitivity of the ratio is perhaps associated to the rate of metabolic activity and thus, early forms and fast-growing stages are more sensitive. For example, Barber *et al.* (1994) used successfully the ratio in 24 hrs exposure of the crustacean *Daphnia magna* to cadmium, and Ibiem and Grant (2005) were also able to use the ratio as a sub lethal endpoint of copper exposure working with a nematode. In this sense, the fish used in the present study are still young fast-growing animals of only 80 days of age (Sellin *et al.*, 2005).

The present study clearly indicates that a short term exposure of copper sulphate, the DNA & RNA content of tissues get reduced in the freshwater fish *N. notopterus* indicating copper sulphate as a pollutant effect the nucleic acid content in the tissues.

Table 1. Showing 96 hrs mortality rate of *N. notopterus* expressed as percentage exposed to copper sulphate toxicant.

Aquarium No.	No. of test animals <i>N. notopterus</i> fish	Concentration of the toxicant (Mg/L)	96 hours	
			No. of fishes dead	% Mortality
1.	10	Control (No toxicant)	Nil	Nil
2.	10	15	1	10%
3.	10	20	3	30%
4.	10	25	5	50%
5.	10	30	10	100%

Table 2. Showing DNA & RNA content (mg/gm) in different tissues of the freshwater fish *Notopterus notopterus* on exposure to Copper sulphate (CuSO₄) MALE.

Organs	Brain		Liver		Kidney		Gonads	
	DNA	RNA	DNA	RNA	DNA	RNA	DNA	RNA
Control	130 ± 1.87 SE ± 0.76	45.62 ± 0.68 SE ± 0.27	143 ± 2.07 SE ± 0.84	49.33 ± 0.56 SE ± 0.22	83.75 ± 1.25 SE ± 0.51	47.16 ± 1.23 SE ± 0.50	175.83 ± 1.63 SE ± 0.66	55 ± 1.00 SE ± 0.40
CuSO ₄ exposed	133.52 ± 1 ^{NS} SE ± 0.50	43.68 ± 1.25 ^{NS} SE ± 0.50	136.57 ± 1.52*** SE ± 0.60	45.50 ± 1.60** SE ± 0.50	81.26 ± 1.26 ^{NS} SE ± 0.78	44.55 ± 0.63 ^{NS} SE ± 0.50	145 ± 1.0*** SE ± 0.50	53 ± 1.50 ^{NS} SE ± 0.50

* Each value is expressed as Mean, Standard Deviation and Standard Error of six observations

Tukey method used:

* = P < 0.05

** = P < 0.01

*** = P < 0.001

NS = No significant

Fig. 1. Determination of LC_{50} by straight line graphical interpolation.

Fig. 2. Showing DNA & RNA content (mg/gm) in different tissues of the freshwater male fish *Notopterus notopterus* on exposure to Copper sulphate ($CuSO_4$).

4. CONCLUSION

The nucleic acid content in different tissues such as brain, liver, kidney & testis of both control and copper sulphate (CuSO₄) exposed freshwater fish *N. notopterus* has been studied. The following observation is made in both control and copper sulphate (CuSO₄) exposed fish. The testis contain large amount of DNA in comparison to other tissues. The degree of DNA content in control and copper sulphate (CuSO₄) exposed fish testis >liver>brain>kidney. The RNA content also exhibited similar to that of DNA, having higher amount in the testis.

References

- [1] Ajitkumar, Surabhi Vashistha and Parthasarathi (2002). Exposure to Interferon g Decreases Levels and Activity of Key Cell Cycle Proteins Resulting in Severe Growth Arrest of the Human Non-Transformed Cell Line, WISH. *J. Cancer Sci. Ther.*, 3(1): 013-019.
- [2] Barber, W.E., Smith, R.L. and Weingartner, T.J. (1994). Fisheries oceanography of the northeast Chukchi Sea: final report. US Department of the Interior, Minerals Management Service, Anchorage, Alaska, OCS Study MMS-93-0051.
- [3] Benton, E., Kloepper and Sams, P.J. (1994). *Environ. Toxicol. Chem.*, 13:1483-1496.
- [4] Bhattacharya, L. and Pandey, A.K. (1989). Sonatox induced impairment of ovarian activity in an exotic cichlid *Oreochromis mossambicus* (Linn). *Foliamorphol (Progue)*, 37(1): 33-37.
- [5] Doudoroff, P., Aderson, B.G., Burduck, G.E., Galtsaff, P.S., Hart, W.B., Patrick, R., Stronge, E.R., Surber, E.W. and Van horn (1951). Bio-assay for the evaluation of acute toxicity of industrial wastes to fish. *Sew. Ind. Wastes*, 23: 1310-1397.
- [6] Fisher, R. (1963). *Statistical Methods for Research Worker* Ed 6, Edinburgh.
- [7] Forstner, U. and Prosi, F. (1979). Heavy metal pollution in freshwater systems. IN: *Biological aspects of freshwater pollution* (Ed. Ravera, O), Pergamon Press, Oxford, pp. 129-159.
- [8] Ibiam, U. and Grant, A. (2005). RNA/DNA ration as a sub lethal endpoint for large-scale toxicity with the nematode *Caenorbditis elegans*. *Environ., Toxicol., Chem.*, 24:1155-1159.
- [9] Kim, S., Kim, J. Kang (2004). Physiological and cellular responses in kidney of rockfish, *Sebastes schlegli* fed dietary cadmium. *J. Fish. Sci. Tech.*
- [10] Miliou, H., Zaboukas, N. and Moraitou-Apostolopoulou, N. (1998). Biochemical composition, growth and survival of the Guppy, *Poecilia reticulata*, during chronic sublethal exposure to cadmium. *Archives of Environmental Contamination and Toxicology*, 35: 58-63.
- [11] Rajnarayan, R. and Satyanesan, A.G. (1986). Effect of mercurial fungicide on the gonadal development of the teleostean fish, *Channa punctatus* (B1). *Ecotoxical. Environ. Saf.*, 11: 352-355.

- [12] Sahai, S. (1989). Toxicology effect of some pesticides on ovaries of *Puntitus ticto* in, Trend in Environmental pollution and pesticides toxicology (Eds: Sashikant et al.). Today and Tomorrow, New Delhi, pp. 21-29.
- [13] Sellin, M.K., Tate-Boldt, E. and Kolok, A.S. (2005). Acclimation to Cu in fathead minnows: Does age influence the response? *Aquat. Toxicol.*, 74:97-109.
- [14] Suedecur, C.W. (1946). Statistical methods low a state college press, Ames Iowa.
- [15] Vidal MA, Kilroy GE, Johnson JR, et al. 2006. Cell growth characteristics and differentiation frequency of adherent equine bone marrow-derived mesenchymal stromal cells: Adipogenic and osteogenic capacity. *Vet Surg*, 35:601–10.
- [16] Vineetashukla, Pratima Rathi and Sastry, K.V. (2002). Effect of cadmium individually and in combination with other metals on the nutritive value of freshwater fish *Channa punctatus*, pp. 158-163.
- [17] Wanganeo, A., S Pani, M.J.Nandan, and I.V.Suresh 1994. Incidence in fish mortality in lower lake of Bopal. *Giobios*, 21:145-146.
- [18] Wilder, I.B. and Stanley, J.B. (1989). DNA- RNA ratios an index to growth to Salmonid fishes in laboratory and stream contaminated with carbaryl. *J. Fish Biol.*, 22:165-172.

(Received 17 January 2015; accepted 26 January 2015)