A comparison of phytotoxic potential among the crude extracts from *Parthenium hysterophorus* L. extracted with solvents of increasing polarity

Ujjal Kumar Pati*, Ashim Chowdhury

Department of Agricultural Chemistry & Soil Science, IAS, University of Calcutta, 35-B.C. Road, Kolkata - 700019, India

*E-mail address: ujjalkrpati@gmail.com

ABSTRACT

There is a worldwide search for the safe, effective and eco-friendly compounds of plant origin to combat the weed species and other pests which are responsible for the great impact on the growth and productivity of agricultural crops. In this present study, a comparison was made to evaluate the phytotoxicity potential of sequentially extracted solvent (hexane, ethyl-acetate, methanol) extracts of *Parthenium hysterophorus* L. (aerial parts) in vitro through bench-top seed germination assay (*Vigna radiata* L.). One-way analysis of variance (ANOVA) followed by Duncan's multiple range test (DMRT) were done for statistical analysis of the data. The study reveals that germination, growth and vigour was significantly ($P < 0.05$) reduced by ethyl-acetate and methanol extracts. The present study concluded that phytotoxicity of ethylacetate and methanolic crude extracts of *Parthenium hysterophorus* could be exploited as potential bioherbicide for future weed management programme and the development of bioherbicide for commercial use.

Keywords: *Parthenium hysterophorus*; phytotoxicity; *Vigna radiata*; bioassay; hexane; ethylacetate and methanol

1. INTRODUCTION

Synthetic chemicals improved crop yields quantitatively and qualitatively but has created undesirable impacts on health, environmental quality and development of resistance and cross resistance. Due to these problems, there is a serious efforts to find alternatives to these chemicals. Among those alternatives, phytochemicals shows a potential approach for the management of noxious and other pests in a proper manner. Besides pest management they are environment friendly, because they are biodegradable, rarely contain halogenated atoms, and possess novel target sites that are different from synthetic chemicals. They also exhibit a large degree of novelty and structural diversity and are highly sought after for the discovery of new agrochemicals. The discovery of novel modes of action would lead to the development of new weed management tools.

There are many weed species that are phytotoxic in nature. Among the weeds *Parthenium hysterophorus* L., is an aggressive weed native to Southern North America, Central America, the West Indies and Central South America[1], having allelopathic effect and drastically retards the growth of many species[2]. *Parthenium* is an invasive exotic weed

commonly known as Bitter weed, False ragweed, Fever few, Ragweed, Carrot weed, White top, Chatak chandani, Congress grass & Star weed. Its botanical name is *Parthenium hysterophorous* L. and belongs to the Class: Magnoliopsida, Family: Asteraceae (Compositae). It is a herbaceous plant, and a native of Tropical America. It is an annual herb and has a deep taproot and erect stem, which becomes woody with age. Height varies of *Parthenium* weed between 50-150 cm, leaves are deeply lobed. It is pale green in colour and has soft hair. *Parthenium* weed flower is creamy white in color. The weed has a large number of highly branched stems. It has small (1-2mm long) black seeds with white scales. They are not visible to the naked eye. The countries where it has been reported with the presence of this weed are South Africa, Mozambique, Madagascar in Africa, China, India, Vietnam, Nepal in Asia, Guatemala, Honduras, Belie etc. in Central America, Mexico, in South America and in United States [3].

The successful spread of *Parthenium* in so many parts of the world including India has mainly been attributed to its allelopathic properties, which enables it to compete effectively with crops and pasture species[4]. It can play an important roles in the determination of plant diversity[5], dominance, succession, and climax of natural vegetation and in the plant productivity of agro ecosystems. Apart from its competitive ability, the invasiveness of the noxious weed *P. hysterophorus* L., is thought to be due to an ability to displace other species by means of allelopathy. The allelochemicals released from *Parthenium* affecting many plant species are sesquiterpene lactones and phenolics[6]. Parthenin is the major sesquiterpene lactone whereas caffeic, vanillic, ferulic, chlorogenic and anisic acids are the major phenolics[7,8]. These two synergistically acting groups of allelochemicals significantly decrease the seed germination and subsequent growth in many crops [8].

2. MATERIAL AND METHODS

2.1. Plant material for extraction

The plant sample (*Parthenium hysterophorous* L) was collected from local vegetation of Baruipur, 24-Paraganas (south) [22°21'29''N Latitude and 88°26'37''E Longitude and altitude of 42 feet] , West Bengal, India. The samples were cut into small pieces, air dried at room temperature ($25 \pm 3^{\circ}$ C), then oven dried for 2 days at 40° C and powdered by mechanical grinder to store in air tight packets which are now ready for experiment.

2.2. Seed material for bioassay

Vigna radiata L. Seeds were used in this experiment as target organism for bioassay. The seed materials were procured from the Department of Agronomy, BCKV, Mohanpur, Kalyani, Nadia (W.B.).

2.3. Extraction

The crude extracts were prepared by macerating 250 gm. of powdered *Parthenium* with 1 L. solvents of choice with increasing polarity (hexane, ethyl-acetate and methanol) following exhaustive sequential extraction. Extraction was done in multiple set and pooled together. The crude extracts were filtered through Whatman No.1 filter paper and were evaporated to a gummy mass, which were then diluted with minimum solvents & remaining

with distilled water to get 10%, 25%, 50%, 75% and 100% (v/v) test extracts for further testing.

2.4. Bioassay

10%, 25%, 50%, 75%, 100% (v/v) test extracts thus obtained from hexane, ethyl acetate and methanol crude extracts were used for bioassay along with a treatment as control with distilled water. A commercial herbicide Pendiguard (pendimethylene 50%EC) at FR dose was used as standard check control. Physiological Experiments of the investigation were carried out with viable surface sterilized seeds of *Vigna radiata*, pretreated with the test extracts of different concentrations for 2 hrs, and placed in Petri dishes containing moistened filter paper.

Germination data were recorded every 12 hr. intervals up to 120 hr of seed soaking following the International Rules for Testing[9]. Seven germination indices, germination percentage (GP), germination index (GI), germination energy (GE), speed of emergence (SE), mean germination time (MGT), seedling vigour index (SVI), and coefficient of the rate/velocity of germination (CRG) were obtained & then calculated from the same data by using the equations[10] described below (Table-1). Morphological parameters like root length, shoot length and seedling height were measured in fresh samples after 120 hrs. Phytotoxicity, inhibition and relative root elongation were calculated following the equations[11] described. One-way analysis of variance (ANOVA) followed by Duncan's multiple range test (DMRT) were done for statistical analysis of the data.

Table 1. Equations used to calculate Germination indices and other physiological parameters.

Germination parameters	Equation
Germination percent (GP)	$GP = [(\text{number of germinated seeds at final count}) / \text{total number of seeds sets for bioassay}] \times 100$
Germination index (GI)	$GI = \sum G_T / T_t$, G_T = number of germinated seeds, T_t = time in days
Mean germination time (MGT)	$MGT = [\sum (T_i \times N_i) / \sum N_i]$ N_i = number of newly germinated seeds at time T_i .
Seedling vigour index (SVI)	$SVI = (\text{seedling length} \times \text{germination percent}) / 100$
Speed of emergence (SE)	$SE = (\text{number of germinated seeds at the starting day of germination} / \text{number of germinated seeds at the final day of measurements}) \times 100$
Germination energy (GE)	$GE = (\text{percent of germinated seeds at the day of germination} / \text{total number of seeds set for bioassay}) \times 100$
Coefficient of the rate of germination (CRG)	$CRG = [(N_1 + N_2 + \dots + N_n) / (N_1 \times T_1 + N_2 \times T_2 + \dots + N_n \times T_n)] \times 100$ where N_1 = number of germinated seeds at time T_1 , N_2 = number of germinated seeds at time T_2 , N_n = number of germinated seeds at time T_n .
Phytotoxicity	$(\text{Root length in control} - \text{Root length in treatment}) / \text{Root length in control}$
Inhibition	$100 - (E_2 \times 100 / E_1)$, Where I = %Inhibition, E_1 = Response in control, E_2 = Response in treatment plant.
Relative root Elongation	$R = (L / L_r) \times 100$, where R is the relative root elongation, L is the mean root length in treatments, L_r is mean root length the of control.

3. RESULTS

The hexane, ethyl acetate and methanol crude extracts of *Parthenium* were used to study the germination process of *V. radiata* seeds upto 120 hrs. The Germination Kinetics was studied for the three crude extracts (Figure-1). It has been revealed that there are nonsignificant changes in germination kinetics (Figure-1A) in hexane crude dilutions where as there is a significant changes in germination kinetics among the treatments of ethyl-acetate (Figure-1B) and methanol (Figure-1C) crudes. The ethyl-acetate treatments show a 50% reduction of germination in comparison to control (100% germination). Table-2-4 shows the effect of crude extract dilutions on germination indices.

Figure 1. Germination kinetics of germinating *V. radiata* seeds treated with different concentrations of crude extracts of *Parthenium* (A- hexane; B- ethylacetate; C- methanol crudes).

Table 2. Effect of hexane crude extract dilutions of *Parthenium* on germination.

TREATMENTS	GP	GI	MGT	CRG	SE	GE	SVI
10	96.67 ^{ab}	94.11 ^{abc}	1.06 ^d	94.22 ^{ab}	95.06 ^a	96.67 ^{ab}	18.49 ^{cd}
25	87.67 ^{bc}	83.33 ^{cd}	1.11 ^d	90.32 ^{ab}	90.07 ^a	87.67 ^{bc}	22.45 ^{bcd}
50	79.33 ^c	73.78 ^d	1.17 ^{cd}	85.45 ^{bc}	86.85 ^a	79.33 ^c	18.62 ^{cd}
75	88.00 ^{bc}	84.34 ^{cd}	1.11 ^d	90.20 ^{ab}	93.27 ^a	88.00 ^{bc}	18.81 ^{cd}
100	84.67 ^c	72.00 ^{bcd}	1.32 ^{bc}	77.10 ^{cd}	70.75 ^b	84.67 ^c	16.78 ^d
CONTROL	100.00 ^a	99.67 ^d	1.01 ^d	99.35 ^a	99.33 ^a	100.00 ^a	31.17 ^{abc}
ST CHK	99.33 ^a	97.83 ^d	1.03 ^d	97.12 ^{ab}	96.98 ^a	99.33 ^a	9.87 ^d
H	98.00 ^a	71.39 ^d	1.70 ^a	58.97 ^e	52.72 ^c	97.83 ^a	33.94 ^{ab}
EA	97.67 ^a	79.81 ^d	1.42 ^b	71.42 ^d	66.26 ^{bc}	97.59 ^a	42.35 ^a
M	99.67 ^a	96.44 ^{ab}	1.07 ^d	93.19 ^{ab}	93.98 ^a	99.63 ^a	38.06 ^a

Data are mean of three replicates; different superscripts in a column indicate significant difference at $P < 0.05$.

Table 3. Effect of ethylacetate crude extract dilutions of *Parthenium* on germination.

TREATMENTS	GP	GI	MGT	CRG	SE	GE	SVI
10	40.67 ^b	26.56 ^d	1.76 ^{ab}	59.07 ^b	39.67 ^{cde}	40.67 ^b	1.15 ^c
25	30.00 ^c	19.00 ^e	1.87 ^a	54.18 ^b	35.51 ^{cde}	30.00 ^c	0.37 ^c
50	14.00 ^d	9.00 ^f	2.02 ^a	50.56 ^b	27.78 ^{de}	14.00 ^d	0.16 ^c
75	2.00 ^e	1.22 ^g	0.67 ^{cd}	16.67 ^c	11.11 ^{ef}	2.00 ^e	0.00 ^c
100	NA	NA	NA	NA	NA	NA	NA
CONTROL	100.00 ^a	100.00 ^a	1.00 ^{bc}	100.00 ^a	100.00 ^a	100.00 ^a	50.53 ^a
ST CHK	100.00 ^a	99.00 ^a	1.02 ^{bc}	98.11 ^a	98.00 ^a	100.00 ^a	34.90 ^b
H	98.00 ^a	71.39 ^c	1.70 ^{ab}	58.97 ^b	52.72 ^{cd}	98.00 ^a	46.17 ^{ab}
EA	97.67 ^a	79.81 ^b	1.42 ^{abc}	71.42 ^b	66.26 ^{bc}	97.67 ^a	40.24 ^{ab}
M	99.67 ^a	96.44 ^a	1.07 ^{bc}	93.19 ^a	93.98 ^{ab}	99.67 ^a	39.82 ^{ab}

Data are mean of three replicates; different superscripts in a column indicate significant difference at $P < 0.05$, NA, not available.

Table 4. Effect of methanol crude extract dilutions of *Parthenium* on germination.

TREATMENTS	GP	GI	MGT	CRG	SE	GE	SVI
10	60.00 ^b	52.11 ^d	1.29 ^{abc}	77.80 ^{bcd}	73.43 ^{ab}	60.00 ^b	6.89 ^c
25	52.67 ^b	44.56 ^{de}	1.38 ^{abc}	72.83 ^{cd}	73.81 ^{ab}	52.67 ^b	3.42 ^c
50	35.33 ^c	29.33 ^{ef}	1.45 ^{abc}	71.42 ^d	67.54 ^{ab}	35.33 ^c	1.76 ^c
75	22.00 ^c	18.22 ^f	1.45 ^{abc}	69.82 ^d	67.10 ^{ab}	22.00 ^c	0.35 ^c
100	34.67 ^c	26.86 ^{ef}	1.79 ^a	65.82 ^d	53.13 ^b	34.67 ^c	0.81 ^c
CONTROL	100.00 ^a	98.61 ^a	1.04 ^{bc}	96.53 ^a	97.67 ^a	100.00 ^a	61.17 ^a
ST CHK	100.00 ^a	99.50 ^a	1.01 ^c	99.03 ^{ab}	99.00 ^a	100.00 ^a	28.37 ^b
H	98.00 ^a	71.39 ^c	1.70 ^{ab}	58.97 ^d	52.72 ^b	98.00 ^a	50.35 ^a
EA	97.67 ^a	79.81 ^{bc}	1.42 ^{abc}	71.42 ^d	66.26 ^{ab}	97.67 ^a	51.27 ^a
M	99.67 ^a	96.44 ^{ab}	1.07 ^{bc}	93.19 ^{abc}	93.98 ^a	99.67 ^a	56.00 ^a

Data are mean of three replicates; different superscripts in a column indicate significant difference at $P < 0.05$

It has been found that all calculated germination indices except MGT, has significantly changed in hexane extract dilutions (Table-2). Ethyl-Acetate treatments at any dilutions (Table-3) have significant ($P < 0.05$) effect on all calculated germination indices except MGT. The GP, GI, CRG, SE, GE, SVI showed a reduction trend with increase in concentration. The 100% ethyl-acetate crude shows least germination followed by 75% treatment which shows maximum reduction as evident by GP(98%) , GI(98.7%) , CRG(83.3%), SE(88.9%), GE(99.3%) and SVI(100%). The lowest concentration (10%) of ethyl-acetate crude treatments also show a significant reduction on all germination indices GP(59.3%), GI(73.4%), CRG(40.93%), SE(60.33%), GE(85.33%) and SVI(97.7%). Almost similar trend of changes were observed in Methanol treatments. At any dilutions (Table-4) significant ($P < 0.05$) effect on all calculated germination indices except MGT were observed. The 75% methanolic treatment shows maximum reduction of germination indices, as evident by GP(78%) , GI(81.5%) , CRG(27.7%), SE(31.3%), GE(84.3%) and SVI(99.4%) as compared to control. On the contrary there is no significant changes for was noted for the H, EA and M solvents as untreated control. The increasing trend of MGT compared to the decreasing trend of GP, GI, CRG, SE, GE, and SVI indicated a significant inhibition or delay of germination of the test species caused by the *Parthenium* crude extracts. The overall result show maximum reduction in ethyl-acetate treatments followed by methanol treatments than the hexane treatments.

Parthenium plant extracts have significant effect on seedling growth of *V. radiata* (Fig- 2).

Figure 2. Effect of *Parthenium* crude extracts on growth (A- hexane; B- ethylacetate; C- methanol crudes).

All the ethyl-acetate treatments (Figure-2B) shows inhibitory effect on the root and shoot growth of the test species followed by methanolic treatments (Figure-2C). There is a decreasing trend in root, shoot and seedling growth as the concentration increases except in the hexane treatments (Figure-2A). At 100% concentration root, shoot and seedling growth was inhibited by 36, 39, and 37% in respect of control respectively for hexane extract; 100, 100 and 100% of control for ethyl-acetate extract; 95, 100 and 96% of control for methanolic extract. The inhibitory effect was greater on the growth of roots than shoot length. The lowest (10%) concentration of ethylacetate and methanol crude extracts also shows a similar trend. The gross toxicity potential of *Parthenium* extracts was corroborated by phytotoxicity and inhibition (Figure-3).

Figure 3. Effect of crude extracts of *Parthenium* on Phytotoxicity and Inhibition of test species (A- hexane; B- ethylacetate; C- methanol crudes).

At 100% concentration of hexane extract (Figure-3A) maximum phytotoxicity and inhibition was observed 37 and 15% respectively. The ethyl-acetate extracts (Figure-3B) and

methanolic extracts (Figure-3C) of all concentration shows significant ($P < 0.05$) toxicity on growth of test species. In both the treatments phytotoxicity and inhibition follows a concentration dependent trend as noted earlier for germination studies. At 100% concentration, phytotoxicity and inhibition was noted by 100 and 100% respectively for ethylacetate extract; 94 and 65 % for methanolic extract, followed by 75, 50, 25, 10% concentrations in both treatments.

4. DISCUSSION

The allelopathic effect of different concentration of solvent extracts of *Parthenium hysterophorus* have been found inhibitory to all parameters viz. Seed germination to metabolism of *Vigna radiata* (mung bean) seeds. Maximum inhibition of germination measured through ‘germination indices’ and growth of *Vigna* seeds were observed for ethylacetate and methanol crude extracts over hexane extracts. Overall the extract at different concentration reduces germinability and slows down germination kinetics which is considered to be the important visible and reliable indices for the evaluation of Allelopathic Effect. GP index indicated the total germination percent of a seed lot after certain period of time when germination became constant. As it is measured by total germination relative to total number of seeds set for germination, GP cannot explain the delayed germination. In contrast, GI is a measure of both percentage and speed of germination and assigns maximum arithmetic weight to seeds that germinate during first count and less weight to those that germinate later. The higher the GI, GE, SE, SVI, and CRG values compared to control, the lower the inhibition, and vice versa, except for MGT [10]. The delay or inhibition of germination caused by phytotoxic plant extracts or substances was also reported by Anjum [12] and Hussain [13] as also corroborated by Tefera [2], Regina et al, [14].

5. CONCLUSION

Therefore in conclusion, ethylacetate and methanolic crudes *Parthenium* extracts has a strong inhibitory effect on seed germination. However the effect is more prominent in ethylacetate crude extracts over methanol extracts. Hence it could be a potential source of bioherbicide with proper management. Since *Parthenium* has become a major invasive in different parts of the country. Therefore all efforts should be made to manage its use in bioherbicide formulation which may be beneficial to farmers as well as environment friendly.

References

- [1] J. Picman, and A.K. Picman, *Biochemical Systematics and Ecology* 12 (1984) 287-297.
- [2] T. Tefera, *Journal of Agronomy and Crop Science* 188 (2002) 306-310.
- [3] R. K. Kohli and D. Rani D, *Research Bulletin (Science), Punjab University* 44 (1984) 105-149.
- [4] R.K. Kohli, D. Rani, H.P. Singh, J.K. Pandher , *Weed Biology and Management* 5 (2005) 105-109.
- [5] C.H. Chou, Y.F. Lee, *J. Chem. Ecol.* 17 (1991) 2267-2281.

-
- [6] C. Swaminathan, R.R.S. Vinaya, K.K. Sureshi, *Int. Tree Crops J.* 6 (1990) 143-150.
- [7] S.D. Kanchan, Jayachandra, *Plant and Soil* 55 (1980) 67-75.
- [8] D.R. Batish, H.P. Singh, R.K. Kohli, S. Kaur, D.B. Saxena, S. Yadav, *Zeitschrift für Naturforschung* 62c (2007) 367-372.
- [9] International Seed Testing Association, *Seed Science Technology* 4 (1976) 51-177.
- [10] A.K.M.M. Islam, H. Kato-Noguchi, <http://dx.doi.org/10.1155/2014/676242>.
- [11] B.J. Rho, B.S. Kil, *J. Nat. Sci. Wankwang Univ* 5 (1986) 19-27.
- [12] T. Anjum, B. Bajwa, *International Journal of Agriculture and Biology* 7 (2005) 417-419.
- [13] M.I. Hussain, L. Gonzalez-Rodriguez, M.J. Reigosa, *Allelopathy Journal* (22)1 (2008) 101-110.
- [14] G.B. Regina, C.F. Reinhardt, L.C. Foxcroft, K. Hurlea, *Crop Protection*, 26 (2007) 237-245.

(Received 21 January 2015; accepted 26 January 2015)