Effect of ethrel on the starch sugar changes of off-season fruits of mango (*Mangifera indica* L. Var. Neelum) during ripening

T. Venkatesan*, C. Tamilmani

Department of Botany, Annamalai University, Annamalai Nagar, Tamil Nadu, India

*E-mail address: dr_venkatesan75@yahoo.com

ABSTRACT

The present investigation is aimed at studying the effect of ethrel on the ripening of off-season fruits of *Mangifera indica* L. var. Neelum. The control fruits were kept in the laboratory naturally while the experimental fruits were treated with different concentrations of ethrel (100, 200 and 300 ppm). In control fruits, partial ripening led to incomplete metabolic changes, which did not alter the presence of sourness in the fruits. Hence, they were not fit to be eaten. On the other hand, the fruits treated with different concentrations (100, 200 and 300 ppm) of ethrel ripened on 13th day, 11th day and 9th day respectively after treatment. The colour changed from green greenish to yellow and the fruits were palatable in nature. The starch decreased during ripening, both in the treated and control fruits. On the other hand, the sugar, α -amylase, β -amylase, activities increased. Among the different 100, 200 and 300 ppm ethrel treatments, the 200 ppm alone had the optimum effect on the ripening of off-season fruits of *Mangifera indica* L. var. Neelum.

Keywords: Starch; Sugar; amylase and fruit ripening

1. INTRODUCTION

Ripening in mangoes involves numerous metabolic activities leading to changes in carbohydrates and acids resulting in declined sugar, acid ratio and development of colour, flavour characteristics, and softening of the texture of acceptable quality. These changes take place in harvested fruits, within a short period of 9 to 12 days at ambient temperature, depending on the variety and stage of maturity. Each mango variety on ripening has distinct characteristics and flavour. Unlike unripe fruits which are astringent, acidic and rich in vitamin C, the ripe fruits are sour and/ or sweet, rich in carotenoids, moderate in vitamin C and highly aromatic (Selvaraj, 1993). As fruits begin to soften, starch deposits are degraded and sugar and flavour components are accumulated (Bathgate *et al.*, 1985). Hydrolysis of starch is a major event during ripening of fruits (Loesecke and Von, 1949). The breakdown of starch to glucose, fructose or sucrose is a characteristic of ripening event. There are several

enzymes in plant tissue that are capable of metabolizing starch (Presis and Levi, 1980). Jain *et al.* (2003) observed that Guava Starch, which is the main storage polysaccharide in many unripe fruits, is degraded during ripening, resulting in sweetness and textural changes in fruits. Guava fruits also exhibited a decrease in starch and an increase in the content of reducing and non-reducing sugars during ripening. Starch content decreased significantly from 3.42 % at MG stage to 0.90 % at overripe stage. Starch content has also been reported to decrease in fruits like papaya. Thanaraj *et al.*, (2009) observed that in Srilankan mango starch concentration was significantly higher in CV. Malgova (35 % DW) than CVS. Karutha colomban (29 % DW) and Willard (21 % DW) fruit. Even though sugar concentration was relatively low in CV. Karutha colomban as compared to CV. Willard, starch concentration was high. Starch levels varied significantly between fully mature (38.6 % DW) and immature stages (19.7% DW). Mango peel (24.4 % DW) had a significantly lower concentration of starch than pulp tissue (30.2 % DW); however, there was no noticeable difference in starch concentration between peel and pulp at immature stage. It was also noticed that the outer pulp (31.6 % DW) contained more starch than inner pulp (28.7 % DW). Dry mature as a proportion of FW increased with maturity and was significantly higher in peel (24.5-0.7 g/100g FW) than in pulp (16.5-31.1 g/100g FW) tissue. Mango CV. Willard had significantly lower dry matter than that of other cultivars.

The flavour of a fruit is compounded mainly of its content of sugars, acids and of numerous volatile aroma components, which are present in very small quantities but which elicit a considerable olfactory response. Changes of flavour during post-harvest ripening typically result from an increase in sugar at the expense of reserve carbohydrate, a decrease in acids, which may be respired, and considerable increase in the production of volatile aroma components (Burton, 1982). The content of sugar increase during fruit ripening was studied in detail by a number of workers (Paul and Southgate, 1978; Tandon and Kalra, 1983; Tandon *et al.*, 1985 Tucker, 1993; Yamaki, 1995; Venkitakrishnan *et al.*, 1997; Lester *et al.*, 1998).

In general, after harvest the sugar content increases at the expense of starch. At the commencement of ripening, the majority of sugars are reducing in nature, but the ripe fruit contains more non-reducing than reducing sugars. In mango, mostly glucose and fructose are the reducing sugars and sucrose is the non-reducing sugar (Subramanyam *et al.*, 1972). A decline in starch content and amylase activity was reported after 91 days of growth in Dasher variety. Glucose and fructose were more until maturity whereas during ripening, sucrose was more plentiful (Tandon and Kalra, 1983; Kalra and Tandon, 1983). In Langra and Mailika, the total sugars, reducing sugars and fructose increased and starch decreased during ripening (Tandon *et al.*, 1985). Sucrose, glucose and fructose were the main sugars in seven mango varieties, and their concentration increased during ripening. Sucrose content which was low at harvest maturity, increased considerably during ripening and was at a comparable level with that of fructose in the varieties, Alphonso and Suvarnakha, and with that of glucose in Banganapalli. Changes in glucose: fructose ratio during ripening showed marked differences between varieties. It increased in the varieties, Suvarnakha and Totapari, decreased in Alphonso, Fazli, and Dasher and remained more or less the same in Banganapalli and Langra (Selvaraj *et al.*, 1989). Thanaraj *et al.* (2009) observed that in Srilankan mango, fructose, glucose and sucrose varied significantly among cultivars. Fructose was the dominant sugar (63.7-130 mg/g DW) in all cultivars and contributed to more than half of total sugar present, followed by glucose (18.6-83.6 mg/g DW) and sucrose (19.8-50.5 mg/g DW). Total sugar was highest in CV. Malgova (260 mg/g DW) followed by CV. Willard (205 mg/g DW) and CV. Ampalavi (190 mg/g DW). There was no significant variation in sugar concentration

according to vertical sectioning (Stem end, middle and distal end). Sugar concentration was significantly lower in peel (119 mg/g DW) than in pulp (202 mg/g DW). However, there was no significant variation between inner and outer pulp. In general, total sugar concentration declined significantly from immature stage (199 mg/g DW) to fully mature stage (162 mg/g DW) of mango. However, sugar concentration was relatively high in the fully mature stage versus the immature of mango CVS. Willard and Ampalavi fruit. The total sugars increased from 4.76 % at MG stage to 8.96 % at OR stage. Both reducing and non-reducing sugars were present at the same concentration (2.38 %) at MG stage. Reducing sugars increased substantially to 5.60 % at OR stage, whereas non-reducing sugars increased slightly and that also at later stages only, i.e., between ripe and overripe stages. This increase is mainly due to degradation of starch.

Sacher (1973) has reported an increase in the activity of amylase during ripening of tomato and mango. An increased amylase activity in banana during ripening was noticed by (Mao and Kinsella, 1981). In Banana, the relative activities of hydrolytic and oxidative enzymes, Viz., α -amylase, starch phosphorylase, acid phosphorylase, Catalase and peroxidase increased in Banana varieties, Pachabale, Rasabale, and Rajabale during ripening. An upsurge in the activities of all the enzymes to a maximum of 1.2 to 19.1 times of the initial level was observed (Desai and Deshpande, 1978). The starch that has accumulated in the maturing fruit is rapidly lost during ripening (Selvaraj *et al.*, 1989) and this loss is evident in the chloroplast where the starch granules become progressively smaller as ripening proceeds. Starch granules completely disappear in the ripe fruit (Parikh *et al.*, 1990) which usually contains negligible levels of starch (Fuchs *et al.*, 1980).

Starch hydrolysis in the ripening mango has been associated with amylase activity (Fuchs *et al.*, 1980), which exhibits the properties of both α and β -amylases. The complete disappearance of starch may be attributed to an upsurge of amylase as ripening is completed.

As a consequence of starch hydrolysis, total sugars increase during ripening, with glucose, fructose and sucrose constituting most of the monosaccharides (Selvaraj *et al.*, 1989). The total sugar content of the ripe „Carabao“ mango is one of the highest reported, with values exceeding 20 % (Peacock and Brown, 1984). However, the lower sugar contents reported for other varieties such as „Golek“ (Lam *et al.*, 1982) might simply reflect differences in the degree of ripeness when optimum eating quality is attained.

α -Amylase and β -amylase are the two amylases in plant tissues capable of metabolizing starch, α -Amylase hydrolyze the α -1,4-linkages of amylose at random to produce a mixture of glucose and maltose, whereas β -amylase attacks only the penultimate linkage from the non-reducing end and thus release only maltose. These enzymes are unable to degrade the β -(1-6) branch points of amylopectin, which are catalyzed by debranching enzymes. Amylase activity increases to some extent during ripening of many fruits (Fuchs *et al.*, 1980; Tucker and Grierson, 1987). Mango and banana are the major starch containing fruits (-15 to 20 %, on fresh weight basis), where starch is almost completely hydrolyzed to free sugars, thus contributing to loosening of structure and textural softening during ripening by Jain *et al.* (2003) observed that Guava the activity of α and β -amylase decreased throughout the process of ripening, suggesting that perhaps the activity of these enzymes in raw guava was sufficient to hydrolyze starch during ripening which was apparent from the decrease in starch content during ripening. The highest activity of starch-hydrolyzing enzymes was associated with the highest starch concentration. The main objective of this was to look the effect of ethrel on the starch, sugar conversion and hydrolytic enzyme activity during the ripening of off-season fruits of mango.

2. MATERIAL AND METHODS

The detached fruits of *Mangifera indica* L. var. Neelum were selected for the present study. The off-season (September to February) green mature unripe fruits were harvested from Auroville near Pondicherry union territory, India and stored in cortons in the Department of Botany at room temperature 28 ± 2 °C with the relative humidity of 85 per cent. They were treated with different concentrations of ethrel (100, 200 and 300 ppm). All the experiments were conducted with seven replicates. The peel and pulp of the fruit material were used to study the ripening process. Starch was extracted and estimated, using the method of Clegg (1956). The residue left behind after the alcoholic extract of the material was taken for starch extraction and estimation. Starch was solubilized with 52 per cent perchloric acid for 50 minutes, filtered, and was made upto 100 ml in a volumetric flask, with distilled water. One to two ml of the perchloric acid extract was diluted with 5 ml of deionised water in test tube and 10 ml of anthrone reagent was added in cold. The contents were heated for 7.5 minutes at 100 °C in a boiling water bath. The test tubes were cooled rapidly and the colour intensity was read at 630 nm in a Spectronic - 20. The starch content was calculated, using a standard graph prepared with glucose. Soluble sugars, reducing and non-reducing sugars, were estimated following the method of Nelson (1944). Two g of fruit material was macerated in a mortar and pestle with 80 per cent ethyl alcohol. The homogenate was centrifuged at 800 rpm for 15 minutes. The supernatant was saved and made upto 20 ml with 80 per cent ethyl alcohol. This extract was used to estimate both reducing and non-reducing sugars. To 1 ml of ethanolic extract, 1 ml of fresh Nelson's reagent (prepared by mixing copper tartrate solution and copper sulphate solution 25:1 (v/v)) was added. The mixture was heated in boiling water for 20 minutes, and then cooled. To the cooled mixture, 1 ml of Nelson's Arsenomolybdate reagent was added. The solution was diluted to 25 ml with distilled water. The intensity of the resulting blue colour was read at 520 nm in a Spectronic -20. The content of the reducing sugar was calculated from glucose standard graph.

Non-reducing sugars were hydrolysed to reducing sugars, and the total sugar was estimated. One ml of ethanolic extract was evaporated to dryness in a boiling water bath. To the residue, 1 ml of distilled water and 1 ml of concentrated H_2SO_4 were added. The mixture was hydrolysed by incubating in an oven at 50 °C for 30 minutes. The solution was neutralized with 1 N NaOH. Total sugar of the hydrolyzed sample was estimated by using Nelson's Arsenomolybdate method. Non-reducing sugar content was calculated by subtracting the value of reducing sugar from the total sugar.

α -amylase and β -amylase activities were assayed, using the modified method of Danielson (1947) and Englard and Singer (1950). Enzyme extraction 1 g of the fruit material was homogenized in a prechilled mortar and pestle with 20 ml of distilled water. The homogenate was centrifuged at 24,000 rpm for 30 minutes at 4 °C in a refrigerated centrifuge. The supernatant was saved and it was used as an enzyme source. To the 0.5 ml of the enzyme extract, 1 ml of 0.1 M citrate buffer (pH 5.0) and 0.5 ml of 2 per cent soluble starch was added. The reaction was allowed for 5 minutes after the addition of starch at 30 °C. After 5 minutes, the reaction was stopped by adding 2 ml of colour reagent. The mixture was boiled for 5 minutes in a water bath at 50 °C. After cooling, the final volume of the solution was made upto 10 ml with distilled water. The absorbance was read at 540 nm in a Spectronic-20. α -amylase activity was assayed by adding 1 ml of 0.1 M citrate buffer (pH 3.4) and 0.5 ml of 2 per cent starch to the 0.5 ml of the enzyme extract. The reaction was allowed for 5 minutes. Then the reaction was stopped by adding

2 ml of the colour reagent and the final volume was made upto 10 ml with distilled water. The colour intensity was read at 640 nm in Spectronic –20.

3. RESULTS AND DISCUSSION

The Table 1 shows the changes in starch content, which occur during the ripening of *Mangifera indica* fruits. The total starch content gradually decreased in the treated fruits, while control fruits retained certain amount of starch during the course of ripening. The starch content was more in control than in treated fruits.

Table 1. Effect of ethrel on the starch changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum.

(Values are Mean $\pm$ SE of 7 samples expressed in mg. Glucose equivalent/g fr. wt.)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE
1	0.115 $\pm$ 0.009	0.108 $\pm$ 0.008	0.102 $\pm$ 0.008	0.090 $\pm$ 0.007	0.105 $\pm$ 0.008	0.096 $\pm$ 0.008	0.092 $\pm$ 0.007	0.080 $\pm$ 0.006
3	0.095 $\pm$ 0.007	0.078 $\pm$ 0.005	0.086 $\pm$ 0.006	0.072 $\pm$ 0.005	0.086 $\pm$ 0.006	0.067 $\pm$ 0.005	0.067 $\pm$ 0.005	0.063 $\pm$ 0.004
5	0.086 $\pm$ 0.005	0.074 $\pm$ 0.004	0.076 $\pm$ 0.005	0.069 $\pm$ 0.004	0.074 $\pm$ 0.004	0.062 $\pm$ 0.004	0.062 $\pm$ 0.004	0.058 $\pm$ 0.003
7	0.075 $\pm$ 0.004	0.070 $\pm$ 0.004	0.062 $\pm$ 0.003	0.066 $\pm$ 0.003	0.064 $\pm$ 0.003	0.056 $\pm$ 0.003	0.050 $\pm$ 0.003	0.054 $\pm$ 0.003
9	0.070 $\pm$ 0.005	0.066 $\pm$ 0.005	0.056 $\pm$ 0.004	0.062 $\pm$ 0.004	0.062 $\pm$ 0.005	0.054 $\pm$ 0.004	0.048 $\pm$ 0.004	0.050 $\pm$ 0.004
11	0.066 $\pm$ 0.005	0.062 $\pm$ 0.004	0.048 $\pm$ 0.003	0.056 $\pm$ 0.004	0.058 $\pm$ 0.004	0.052 $\pm$ 0.004	0.036 $\pm$ 0.003	0.048 $\pm$ 0.003
13	0.058 $\pm$ 0.003	0.052 $\pm$ 0.003	0.036 $\pm$ 0.002	0.048 $\pm$ 0.003	0.049 $\pm$ 0.003	0.046 $\pm$ 0.003	0.028 $\pm$ 0.002	0.034 $\pm$ 0.002
15	0.046 $\pm$ 0.002	0.042 $\pm$ 0.002	0.028 $\pm$ 0.001	0.036 $\pm$ 0.002	0.038 $\pm$ 0.002	0.034 $\pm$ 0.002	0.018 $\pm$ 0.001	0.026 $\pm$ 0.001

The content of starch decreased was more in the pulp than in the peel, both in the treated and control fruits. The percentage of loss was more in the 200 ppm ethrel treated fruits than in the 100, 300 ppm and control. As fruits began to soften, starch deposits degraded and sugar and flavour components accumulated (Bathgate *et al.*, 1985). Hydrolysis of starch was a major event during the ripening of fruits (Loesecke and Von 1949). The breakdown of starch to glucose, fructose or sucrose, is a characteristic of ripening event. There are several enzymes in plant tissue, capable of metabolizing starch (Presis and Levi, 1980). In banana, the green and unripe fruits are rich in reserve

carbohydrate in the form of starch. During ripening, almost entire starch is converted into simple sugars, such as sucrose, fructose and glucose. Only 1-2 per cent of starch remains in the ripe fruits similarly certain amount of starch remains in the ripened off-season fruit of *Mangifera indica*. The retained starch content was higher in the control fruits than in the treated fruits. Hence, the untreated fruits are not suitable for edible purposes because of high sourness and rich amount of starch. On the other hand, hydrolysis of starch is more in the treated fruits; comparably the content of sugar also increases. Hence it is suitable for edible purposes. Jain et al. (2003) observed that guava starch, which is the main storage polysaccharide in many unripe fruits, is degraded during ripening, resulting in sweetness and textural changes in fruits. Guava fruits also exhibited a decrease in starch and an increase in the content of reducing and non-reducing sugars during ripening. Starch content decreased significantly from 3.42 % at MG stage to 0.90 % at overripe stage. Thanaraj et al. (2009) observed that Srilankan mangoes starch concentration was significantly higher in CV. Malgova (35 % DW) than in CV. Karutha colomban (29 % DW) and Willard (21 % DW) fruits. Even though sugar concentration was relatively low in CV. Karutha colomban as compared to CV. Willard, starch concentration was high. Starch levels varied significantly between fully mature (38.6 % DW) and immature stages (19.7 % DW). Mango peel (24.4 % DW) had a significantly lower concentration of starch than pulp tissue (30.2 % DW); however, there was no noticeable difference in starch concentration between peel and pulp at immature stage. It was also noticed that the outer pulp (31.6 % DW) contained more starch than inner pulp (28.7 % DW). Dry matter as a proportion of FW increased with maturity and was significantly higher in peel (24.5-50.7 g/100g FW) than in pulp (16.5-31.1 g/100g FW) tissue. Mango CV. Willard had significantly lower dry matter than that of other cultivars. The Tables 2, 2(a) and 2(b) show the changes in sugar, which occur during the ripening of *Mangifera indica* fruits.

Table 2. Effect of ethrel on the reducing sugar changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum.

(Values are Mean $\pm$ SE of 7 samples expressed in mg. Glucose equivalent/g fr. wt.)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
1	0.338 $\pm$ 0.027	0.330 $\pm$ 0.026	0.335 $\pm$ 0.027	0.329 $\pm$ 0.026	0.365 $\pm$ 0.029	0.359 $\pm$ 0.029	0.355 $\pm$ 0.028	0.372 $\pm$ 0.030
3	0.344 $\pm$ 0.024	0.338 $\pm$ 0.024	0.340 $\pm$ 0.024	0.336 $\pm$ 0.023	0.370 $\pm$ 0.026	0.369 $\pm$ 0.026	0.358 $\pm$ 0.025	0.376 $\pm$ 0.026
5	0.352 $\pm$ 0.021	0.343 $\pm$ 0.021	0.349 $\pm$ 0.021	0.341 $\pm$ 0.020	0.380 $\pm$ 0.023	0.349 $\pm$ 0.021	0.343 $\pm$ 0.021	0.386 $\pm$ 0.023
7	0.368 $\pm$ 0.018	0.366 $\pm$ 0.018	0.370 $\pm$ 0.019	0.364 $\pm$ 0.018	0.402 $\pm$ 0.020	0.396 $\pm$ 0.020	0.390 $\pm$ 0.020	0.406 $\pm$ 0.020
9	0.386 $\pm$ 0.031	0.390 $\pm$ 0.031	0.393 $\pm$ 0.031	0.389 $\pm$ 0.031	0.429 $\pm$ 0.034	0.424 $\pm$ 0.034	0.410 $\pm$ 0.033	0.430 $\pm$ 0.034
11	0.410 $\pm$ 0.029	0.415 $\pm$ 0.029	0.419 $\pm$ 0.029	0.409 $\pm$ 0.029	0.449 $\pm$ 0.031	0.438 $\pm$ 0.031	0.435 $\pm$ 0.030	0.452 $\pm$ 0.032
13	0.426 $\pm$ 0.026	0.420 $\pm$ 0.025	0.426 $\pm$ 0.026	0.411 $\pm$ 0.025	0.450 $\pm$ 0.027	0.449 $\pm$ 0.027	0.445 $\pm$ 0.027	0.452 $\pm$ 0.027
15	0.442 $\pm$ 0.022	0.426 $\pm$ 0.021	0.432 $\pm$ 0.022	0.419 $\pm$ 0.021	0.468 $\pm$ 0.024	0.459 $\pm$ 0.023	0.448 $\pm$ 0.022	0.470 $\pm$ 0.024

Table 2(a). Effect of ethrel on the non-reducing sugar changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum

(Values are Mean $\pm$ SE of 7 samples expressed in mg. Glucose equivalent/g fr. wt.)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
1	0.342 ± 0.027	0.338 ± 0.027	0.349 ± 0.028	0.336 ± 0.027	0.378 ± 0.030	0.380 ± 0.030	0.376 ± 0.030	0.385 ± 0.031
3	0.348 ± 0.024	0.346 ± 0.024	0.352 ± 0.025	0.344 ± 0.024	0.384 ± 0.027	0.372 ± 0.026	0.380 ± 0.027	0.382 ± 0.027
5	0.359 ± 0.021	0.356 ± 0.021	0.360 ± 0.022	0.348 ± 0.021	0.400 ± 0.024	0.396 ± 0.024	0.392 ± 0.024	0.398 ± 0.024
7	0.380 ± 0.019	0.378 ± 0.019	0.382 ± 0.019	0.372 ± 0.019	0.428 ± 0.022	0.412 ± 0.020	0.412 ± 0.020	0.418 ± 0.021
9	0.398 ± 0.032	0.397 ± 0.032	0.408 ± 0.033	0.394 ± 0.031	0.442 ± 0.036	0.428 ± 0.034	0.424 ± 0.034	0.435 ± 0.035
11	0.425 ± 0.030	0.426 ± 0.029	0.439 ± 0.030	0.421 ± 0.029	0.466 ± 0.032	0.455 ± 0.032	0.451 ± 0.032	0.460 ± 0.033
13	0.436 ± 0.026	0.428 ± 0.026	0.444 ± 0.026	0.422 ± 0.025	0.482 ± 0.029	0.465 ± 0.028	0.455 ± 0.027	0.472 ± 0.028
15	0.439 ± 0.022	0.432 ± 0.022	0.436 ± 0.022	0.429 ± 0.021	0.486 ± 0.024	0.471 ± 0.024	0.461 ± 0.023	0.478 ± 0.023

Table 2(b). Effect of ethrel on the total sugar changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum

(Values are Mean $\pm$ SE of 7 samples expressed in mg. Glucose equivalent/g fr. wt.)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
1	0.677 ± 0.054	0.668 ± 0.053	0.665 ± 0.053	0.687 ± 0.055	0.731 ± 0.058	0.739 ± 0.059	0.750 ± 0.060	0.750 ± 0.060
3	0.688 ± 0.048	0.684 ± 0.048	0.680 ± 0.047	0.696 ± 0.049	0.738 ± 0.052	0.741 ± 0.052	0.752 ± 0.053	0.760 ± 0.053
5	0.708 ± 0.042	0.699 ± 0.042	0.689 ± 0.041	0.710 ± 0.043	0.735 ± 0.045	0.745 ± 0.045	0.778 ± 0.047	0.786 ± 0.047
7	0.750 ± 0.038	0.744 ± 0.037	0.736 ± 0.037	0.750 ± 0.037	0.802 ± 0.040	0.808 ± 0.040	0.820 ± 0.041	0.834 ± 0.042
9	0.791 ± 0.063	0.787 ± 0.063	0.783 ± 0.062	0.794 ± 0.064	0.834 ± 0.067	0.852 ± 0.068	0.864 ± 0.069	0.872 ± 0.070
11	0.844 ± 0.059	0.841 ± 0.058	0.830 ± 0.058	0.849 ± 0.059	0.886 ± 0.062	0.893 ± 0.063	0.909 ± 0.064	0.918 ± 0.064
13	0.862 ± 0.052	0.848 ± 0.051	0.833 ± 0.050	0.870 ± 0.052	0.900 ± 0.054	0.914 ± 0.055	0.922 ± 0.055	0.934 ± 0.056
15	0.871 ± 0.044	0.858 ± 0.043	0.848 ± 0.042	0.878 ± 0.044	0.929 ± 0.045	0.930 ± 0.047	0.946 ± 0.047	0.956 ± 0.048

The reducing sugar, non-reducing sugar and total sugar content gradually increased during the course of ripening, both in the control and treated fruits. The content of sugar was more in the treated fruits than in the control fruits. The non-reducing sugar content was more than that of reducing sugars in the peel and the pulp of the treated and control fruits. The content of sugar was more in the pulp than in the peel during the ripening process. The statistical analysis on starch and total sugar in the peel and pulp, both in the control and treated, showed a negative correlation. The correlation co-efficient values were - 0.93, - 0.87, - 0.92, - 0.96, - 0.90, - 0.82, - 0.92 and - 0.95. The observed correlation co-efficient values were significant at 1 % level. In general, after harvest, the sugar content increases at the expense of starch. At the commencement of ripening, the majority of sugars are reducing in nature, but the ripe fruit contains more non-reducing than reducing sugars. In mango, mostly glucose and fructose are the reducing sugars and sucrose is the non-reducing sugar (Subramanyam *et al.*, 1972). Similarly Thanaraj (2009) reported that in Srilankan Mango the fructose, glucose and sucrose varied significantly among cultivars. Fructose was the dominant sugar (63.7-130 mg/g DW) in all cultivars and contributed to more than half of total sugar present, followed by glucose (18.6-83.6 mg/g DW) and sucrose (19.8-50.5 mg/g DW). Total sugar was highest in CV. Malgova (260 mg/g DW) followed by CV. Willard (205 mg/g DW) and CV. Ampalavi (190 mg/g DW). There was no significant variation in sugar concentration according to vertical sectioning (Stem end, middle and distal end). Sugar concentration was significantly lower in peel (119 mg/g DW) than in pulp (202 mg/g DW). However, there was no significant variation between inner and outer pulp. In general, total sugar concentration declined significantly from immature stage (199 mg/g DW) to fully mature stage (162 mg/g DW) of mango. However, sugar concentration was relatively high in the fully mature stage versus the immature of mango CVS. Willard and Ampalavi fruit. The total sugars increased from 4.76 % at MG stage to 8.96 % at OR stage. Both reducing and non-reducing sugars were present at the same concentration (2.38 %) at MG stage. Reducing sugars increased substantially to 5.60 % at OR stage, whereas non-reducing sugars increased slightly and that also at later stages only, i.e., between ripe and overripe stages. This increase is mainly due to degradation of starch. A decline in starch content and amylase activity was reported after 91 days of growth in the variety, Dasherri. Glucose and fructose were more until maturity, whereas during ripening, sucrose was more plentiful (Tandon and Kalra, 1983; Kalra and Tandon, 1983). In the Langra and Mailika, the total sugars, reducing sugars and fructose increased and starch decreased during ripening (Tandon *et al.*, 1985). Similar findings were observed in the off-season fruits of *Mangifera indica*. Sucrose, glucose, and fructose were the main sugars in seven mango varieties and their concentration increased during ripening. Sucrose content that was low in the fruits harvested at maturity increased considerably during ripening, and was at a comparable level with that of fructose in the varieties Alphonso and Suvarnarekha and with that of glucose in Banganapalli. Changes in glucose: fructose ratio during ripening showed marked differences between varieties. It increased in the varieties Suvarnarekha and Totapari, decreased in Alphonso, Fazli and Dasherri, and remained more or less the same in Banganapalli and Langra (Selvaraj *et al.*, 1989). In mango, sweetness was more due to the presence of the high amount of non-reducing sugar than that of reducing sugar. As the content of starch decreased, the level of reducing and non-reducing sugars increased. The statistical analysis of results on starch and sugar during ripening showed negative correlation. The correlation co-efficient values are highly significant at 1 per cent level. The increase in the sugar was also coupled with the increase in the activity of the enzyme amylase. The disappearance of starch is one of the most dramatic chemical changes associated with the

ripening of many fruits. Starch breakdown can occur either by phosphorylysis catalysed starch phosphorylase or by hydrolysis catalysed by the enzyme amylase. Both may be involved in senescent sweetening of fruits, for example potato (Burton, 1982). In *Mangifera indica* fruit the activity of α -amylase and β -amylase gradually increased, both in the peel and the pulp of the treated and control fruits. The pulp had more α , β -amylase activity than that of the peel, while the α -amylase activity was more than β -amylase activity during the course of ripening, both in the treated and control fruits. Among the treated fruits, the fruits had more activity of α and β -amylase in the fruits treated with 200ppm ethrel than in the 100, 300 ppm and control. (Tables 3 and 4). Of these two hydrolytic enzymes α -amylase and β -amylase, the α -amylase activity was more than the β -amylase. Hence, the α -amylase is a key enzyme in the starch breakdown of *Mangifera indica* fruit. Similar increase of amylase activity in banana was noticed by Mao and Kinsella (1981).

Table 3. Effect of ethrel on the α - amylase changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum.

(Values are Mean $\pm$ SE of 7 samples expressed in maltose equivalent mg/g fr. wt)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE
1	11.33 $\pm$ 0.906	11.28 $\pm$ 0.902	11.44 $\pm$ 0.915	11.36 $\pm$ 0.909	21.57 $\pm$ 1.726	21.58 $\pm$ 1.726	21.57 $\pm$ 1.726	21.58 $\pm$ 1.726
3	12.58 $\pm$ 0.881	13.16 $\pm$ 0.921	15.56 $\pm$ 1.089	14.36 $\pm$ 1.005	22.16 $\pm$ 1.551	22.26 $\pm$ 1.558	22.38 $\pm$ 1.566	22.45 $\pm$ 1.572
5	13.16 $\pm$ 0.790	14.26 $\pm$ 0.856	19.74 $\pm$ 1.184	15.34 $\pm$ 0.920	23.18 $\pm$ 1.391	23.88 $\pm$ 1.433	23.18 $\pm$ 1.391	24.46 $\pm$ 1.468
7	15.28 $\pm$ 0.764	16.36 $\pm$ 0.818	25.96 $\pm$ 1.298	19.26 $\pm$ 0.963	25.16 $\pm$ 1.258	25.82 $\pm$ 1.291	25.16 $\pm$ 1.258	28.42 $\pm$ 1.421
9	17.32 $\pm$ 1.386	18.21 $\pm$ 1.457	27.42 $\pm$ 2.194	22.24 $\pm$ 1.780	27.81 $\pm$ 2.225	28.16 $\pm$ 2.253	31.66 $\pm$ 2.533	35.78 $\pm$ 2.862
11	20.45 $\pm$ 1.432	20.84 $\pm$ 1.458	30.65 $\pm$ 2.145	23.24 $\pm$ 1.627	30.56 $\pm$ 2.139	33.75 $\pm$ 2.362	41.94 $\pm$ 2.936	36.18 $\pm$ 2.533
13	23.36 $\pm$ 1.401	23.86 $\pm$ 1.431	32.36 $\pm$ 1.942	24.16 $\pm$ 1.449	33.68 $\pm$ 2.021	34.18 $\pm$ 2.050	47.92 $\pm$ 2.876	40.16 $\pm$ 2.409
15	26.46 $\pm$ 1.323	26.76 $\pm$ 1.338	34.14 $\pm$ 1.707	28.46 $\pm$ 1.423	36.54 $\pm$ 1.827	40.26 $\pm$ 2.013	54.14 $\pm$ 2.707	46.28 $\pm$ 2.314

Table 4. Effect of ethrel on the β amylase changes during the ripening of off-season fruits of *Mangifera indica* L. var. Neelum.

(Values are Mean $\pm$ SE of 7 samples expressed in maltose equivalent mg/g fr. wt)

Days	Peel				Pulp			
	Control	100 ppm	200 ppm	300 ppm	Control	100 ppm	200 ppm	300 ppm
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE
1	10.62 $\pm$ 0.849	10.42 $\pm$ 0.834	10.84 $\pm$ 0.867	10.34 $\pm$ 0.827	11.58 $\pm$ 0.926	11.72 $\pm$ 0.938	11.84 $\pm$ 0.947	11.82 $\pm$ 0.946
3	11.82 $\pm$ 0.827	11.32 $\pm$ 0.792	12.56 $\pm$ 0.879	14.40 $\pm$ 1.008	12.23 $\pm$ 0.856	12.26 $\pm$ 0.858	12.78 $\pm$ 0.895	12.60 $\pm$ 0.882
5	12.56 $\pm$ 0.754	12.41 $\pm$ 0.745	14.84 $\pm$ 0.890	13.48 $\pm$ 0.809	15.34 $\pm$ 0.920	16.46 $\pm$ 0.988	20.86 $\pm$ 1.252	19.46 $\pm$ 1.168
7	14.48 $\pm$ 0.724	14.52 $\pm$ 0.726	16.68 $\pm$ 0.834	15.42 $\pm$ 0.771	19.26 $\pm$ 0.963	20.16 $\pm$ 1.008	23.48 $\pm$ 1.174	21.36 $\pm$ 1.068
9	15.86 $\pm$ 1.269	15.82 $\pm$ 1.265	18.86 $\pm$ 1.508	17.56 $\pm$ 1.405	21.18 $\pm$ 1.694	22.46 $\pm$ 1.797	28.56 $\pm$ 2.285	25.48 $\pm$ 2.038
11	17.88 $\pm$ 1.252	16.62 $\pm$ 1.163	20.16 $\pm$ 1.411	19.58 $\pm$ 1.371	24.26 $\pm$ 1.698	25.56 $\pm$ 1.789	33.46 $\pm$ 2.342	28.36 $\pm$ 1.985
13	19.28 $\pm$ 1.157	19.18 $\pm$ 1.151	22.48 $\pm$ 1.349	21.18 $\pm$ 1.271	22.16 $\pm$ 1.330	23.12 $\pm$ 1.387	38.18 $\pm$ 2.291	32.36 $\pm$ 1.942
15	22.58 $\pm$ 1.129	22.46 $\pm$ 1.123	24.88 $\pm$ 1.244	23.16 $\pm$ 1.158	24.18 $\pm$ 1.209	25.42 $\pm$ 1.271	44.82 $\pm$ 2.241	38.76 $\pm$ 1.938

α -Amylase and β -amylase are the two amylases in plant tissues capable of metabolizing starch, α -Amylase hydrolyze the α -1,4-linkages of amylose at random to produce a mixture of glucose and maltose, whereas β -amylase attacks only the penultimate linkage from the non-reducing end and thus releases only maltose. These enzymes are unable to degrade the β - (1-6) branch points of amylopectin, which are catalyzed by debranching enzymes. Amylase activity increases to some extent during ripening of many fruits (Fuchs *et al.*, 1980; Tucker and Grierson, 1987).

Mango and banana are the major starch containing fruits (-15 to 20 %, on fresh weight basis), where starch is almost completely hydrolyzed to free sugars, thus contributing to loosening of the all structure and textural softening during ripening (Bhagyalakshmi *et al.*, 2002). Jain *et al.* (2003) noticed a similar observation in Guava: the activity of α and β -amylase decreased throughout the process of ripening, suggesting that perhaps the activity of these enzymes in raw guava was sufficient to hydrolyze starch during ripening which was

apparent from the decrease in starch content during ripening. The highest activity of starch-hydrolyzing enzymes was associated with the highest starch concentration.

4. CONCLUSION

Among the different 100, 200 and 300 ppm ethrel treatment the 200 ppm alone had the optimum effect on the ripening of off-season fruits of *Mangifera indica* L. var Neelum.

References

- [1] Selvaraj Y. (1993). Fruit ripening in Advances in Horticulture, *Fruit Crops*, (ed. K.L. Chadha and O.P. Pareek). Malhotra Publishing House, New Delhi, Vol.4, Part 4, 1703-1741.
- [2] Bathgate B., Purton M. E., Grierson D. Goodenough P. W., *Planta* 165 (1985) 197-204.
- [3] Loesecke H., Von W. (1949). Bananas, *Interscience*, Publishers, New York. p. 189.
- [4] Presis J., Levi C. (1980). Starch biosynthesis and degradation. In: "*The Biochemistry of Plants*" – A comprehensive treatise. (ed. J. Preiss), Academic Press, vol. 13, 371-424.
- [5] Jain N., Dhawan K., Malhotra S., Singh R., *Pl. Fd. Human Nutr.* 58 (2003) 309-315.
- [6] Thanaraj T., Terry L. A., Bessant C., *J. Fd. Chem.* 112 (2009) 786-794.
- [7] Paul A. A., Southgate D. A. T. (1978). The composition of Foods. HMSO. London, 418.
- [8] Tandon D. K., Kalra S. K., *J. Hort. Sci.* 58(3) (1983) 449-453.
- [9] Tandon D. K., Kalra S. K., Lohani H. C., *Ind. J. Hort.* 42 (1985) 223-228.
- [10] Tucker G. A. (1993). *Introduction in Biochemistry of Fruit Ripening*. (ed. G. Seymour, J. Taylor and G. Tucker,) Chapman & Hall, London, pp. 1-51.
- [11] Yamaki S., *Acta Hort.* 398 (1995) 109-120.
- [12] Venkitakrishnan M., Panjatcharam V., Kumaravelu G., Ramanujam M. P., *Ind. J. Pl. Physiol.* 2(4) (1997) 267-270.
- [13] Lester G., *J. Am. Soc. Hort. Sci.* 23(1) (1998) 126-129.
- [14] Subramanyam H., Narayanamoorthy N. V., Lakshmi-narayana S., Santha K., *Acta Hort.* 24 (1972) 260-264.
- [15] Kalra S. K., Tandon D. K., *Sci. Hort.* 19 (1983) 263-269.
- [16] Selvaraj Y., Kumar R., Pal D. K., *J. Fd. Sci. Tech.* 26 (1989) 308-313.
- [17] Sacher J. A., *Ann. Rev. Pl. Physiol.* 24 (1973) 197-224.
- [18] Mao and Kinsella, *Fd. Sci.* 46 (1981) 1400-1403.
- [19] Desai B. B., Deshpande P. B., *Mysore J. Agri. Sci.* 12 (1978) 193-201.

- [20] Parikh H. R., Nair G. M., Modi, V. V., *Ann. Bot.* 65 (1990) 121-127.
- [21] Fuchs Y., Pesis E., Zauberman G., *Sci. Hort.* 13 (1980) 155-160.
- [22] Peacock B. C., Brown B. I. (1984). *Quality comparison of Several mango varieties*. Proceedings First Australian Mango Research Workshop, pp. 334-339.
- [23] Lam P. F., Ng K. H., Omar D., Talib Y. (1982). Physical, Physiological and Chemical changes of „Golek“ after harvest. Proc. Workshop on Mango and Rambutan. ASEAN, Postharvest Training and Research Center, College, Languna, Philippines, pp. 96-112.
- [24] Tucker G. A., Grierson D. (1987). Fruit ripening. In *The Biochemistry of plants – A comprehensive treatise* (ed. D. D. Davies), Academic Press, vol. 12, 265-318.
- [25] Clegg K. M., *Sci. Fd. Agri.* 7 (1956) 40-44.
- [26] Nelson N., *J. Biol. Chem.* 153 (1944) 375-380.
- [27] Danielson C. E., *Nature* 160 (1947) 889.
- [28] Englard S., Singer T. P., *J. Biochem.* 187 (1950) 213.
- [29] Barnell H. R., *Annals. Bot.* 20 (1941) 607-645.
- [30] Barnell H. R., *Annals. Bot.* 20 (1941) 607-645.
- [31] Bhagyalakshmi N., Prabha T. N., Yashoda H. M., Prasanna V., Jagadeesh B. H., Tharanathan R. N., *Acta Hort.* 575 (2002) 717-724.

(Received 26 November 2013; accepted 30 November 2013)