Determination of Five Mineral Element Contents in Pollen Grains of Different Seedling Date Palm (*Phoenix dactylifera* L.) Male Trees Grown in Fayoum Governorate, Egypt

Samir A. Seif El-Yazal^{1a*}, Mohamed A. Seif El-Yazal²

¹Horticulture Department, Faculty of Agriculture, Fayoum University

²Botany Department, Faculty of Agriculture, Fayoum University, Egypt

Author of Correspondence: sas00@fayoum.edu.eg

Keywords: Male date palm, Zinc, Iron; Manganese, Copper, Magnesium.

Abstract. This experiment was conducted to found the connection between sensible male pollinators and the content from mineral element (Zinc, Iron, Manganese, Copper and Magnesium) in pollen grain of fifty seedling date palm male and compared it with the best selective five seedling date palm males (number 2, 10, 29, 40, 46) are appropriate and promising males for to be utilized in pollinating female date palms and mistreatment in breeding programs in previous study. Results found variations among the categories of pollen and elements. The highest concentration of zinc was found in males' number (2, 10, 40 and 46) which recorded 127,102, 115 and 122 µg/g dry weight respectively as compared to the other male trees. Also the highest concentration of iron was found in male number (40) which recorded 222 µg/g dry weight as compared to the other male trees. Moreover, notes from result excellence male number (10) in manganese concentration which recorded 202 µg/g dry weight as compared to the other male trees. In this concern, pollen grain of date palm male trees contained also, high concentrations of copper and magnesium the concentration ranging from 10 to 30 and 1320 to 1985 µg/g dry weight for copper and magnesium respectively. The best value for copper and magnesium which obtained from the selective male number (46) which recorded 30 and 1985 µg/g dry weight for copper and magnesium respectively as compared to the other male trees. From all the results we can conclude that the date palm pollen grain was as rich source of important minerals, so its suitability as a regular component in plant and human diet.

1. Introduction

Date palm (*Phoenix dactylifera* L.) is one of the oldest cultivated trees in world. In Egypt, Date palm tree is taken into account joined of the foremost vital fruits wherever it's cosmopolitan in numerous areas in Egypt from the Mediterranean coast up to Aswan. It's cosmopolitan in numerous districts of the planet districts of the world. It constitutes the foremost vital plant in arid and desert areas wherever it provides favorable conditions for both human and animal habitats [1]. In fact, palm are irreplaceable tree in irrigable desert lands grows underneath unfavorable conditions; wherever several of the opposite fruit species doesn't. It additionally keeps alternative crops settled under that from heat, wind, and even cold weather. So, it plays an enormous role to prevent geologic process and provides life to desert areas [2, 3]. Date palm is a dioecious perennial and monocotyledonous fruit tree that belongs to the Arecaceae family. Several investigators studied the effect of pollens obtained from totally different palm males cultivar on fruit set, yield and fruit physical and chemical properties of many female palm cultivars [4, 5, 6, 7, 8, 9, 10]. The direct influence of the male parent on the event of the date fruit is precise and definite and varies with the actual male went to fertilize the female flowers. Every male exerting just about constant impact on fruit of all varieties and exerting constant impact in several years. Therefore, it's vital to select and identify superior male in term of fertilization. The concentration of mineral components in pollen grains of date palm males were studied by many workers [11,12,13,14,15,16,17,18,19,20,21]. In addition, several investigators mentioned that, the effect of micro-nutrient in pollen grain germination and pollen tube growth in several plant species [22, 23, 24]. The minerals content in pollen grains differed from one seedling plant males to a different. The

macronutrients, N was gift within the highest concentrations, followed by Ca, P, Mg and Na, whereas the micronutrients, Fe was gift within the highest concentrations, followed by Zn, Mn and Cu [25]. The aim of this study is to determine the content of those five types elements for minerals (Zinc, Iron, Manganese, Copper and Magnesium) in pollen grain of the selective five seedling date palm males and compare them with those of alternative males because these elements are importance within the life of the plant wherever any shortage leads to intensive harm to the plant. Moreover, these males might be used as pollinators for palm females. Also, we are able to propagate and multiply these promising males' oldsters by tissue cultural programs. Moreover, we wish to indicate is that the content of those elements are correlated with the rise in pollination and fertilization process or not. Also, we are able to propagate and multiply these promising males' oldsters by tissue cultural programs. Moreover, wish to indicate is that the content of those elements are correlated with the rise in pollination and fertilization process or not

2. Materials and Methods

2.1. Plant materials

This study was conducted during the three successive seasons 2015, 2016 and 2017 on fifty seedling date palm males about 30 years old and grown in Sand clay loamy soil at Tamiya district, Fayoum Governorate, Egypt., the study included the five date palm males coded by numbers (2, 10, 29, 40 and 46) which select according to the evaluation results of previously studies[8,9,10] as good male pollinators for further using in pollinating female date palms and breeding programs.

2.2. Collection of plant material

Date palm pollen was collected in March (2015, 2016 and 2017) from Tamiya district, Fayoum Governorate, Egypt. Some of palm pollen grains were collected and separated from the kernels with a fine gauze sieve and dried at 70°C and then kept in closed container.

2.3. Chemical composition:

Dried pollen grain were used for determination, iron, zinc, manganese, copper and magnesium contents as $\mu\text{g/g}$ dry weight by using atomic absorption spectrophotometer apparatus according to [26,27].

Data were statistically analyzed according to [28]. Data presented in this investigation represent the mean of the three experimental seasons.

3. Results

3.1. Micro nutrient

3.1.1. Zinc concentrations

Data presented in Table (1) indicate that pollen grain of date palm male trees contained high concentrations of zinc, the concentration ranging from 49 to 127 $\mu\text{g/g}$ dry weight. The best result which obtained from the selective males number (2, 10, 40 and 46) recorded 127, 102, 115 and 122 $\mu\text{g/g}$ dry weight respectively as compared to the other male trees.

Table1. Pollen zinc content of date palm males as the mean of the three seasons

Male No.	Zn µg/g dry weight		Male No.	Zn µg/g dry weight	
	Mean*	C.V.**.		Mean*	C.V.**
1	88	3.13	26	92	1.25
2	127	3.62	27	95	1.95
3	95	1.33	28	70	2.41
4	93	6.31	29	98	2.63
5	74	1.43	30	81	1.01
6	55	2.67	31	63	2.14
7	62	2.61	32	66	1.92
8	81	1.43	33	53	2.33
9	49	3.58	34	84	1.36
10	102	4.14	35	84	1.85
11	73	1.40	36	90	2.06
12	92	6.25	37	77	1.35
13	94	6.71	38	94	1.18
14	82	1.66	39	63	2.55
15	83	1.86	40	115	3.02
16	83	1.17	41	51	1.18
17	60	2.01	42	65	1.66
18	64	1.95	43	83	1.22
19	58	1.18	44	90	1.28
20	79	1.77	45	95	1.35
21	88	2.03	46	122	2.47
22	67	1.18	47	59	1.35
23	72	3.64	48	57	1.48
24	75	3.22	49	92	1.79
25	83	1.84	50	64	1.45

*Mean of seasons, 2015, 2016and 2017

**Coefficient of variation

3.1.2. Iron concentrations

Data presented in Table (2) indicate that pollen grain of date palm male trees contained also high concentrations of iron, the concentration ranging from 145 to 222µg/g dry weight .The best result obtained from the selective male number (40) which recorded 222 µg/g dry weight respectively as compared to the other male trees.

Table 2. Pollen iron content of date palm males as the mean of the three seasons

Male No.	Fe µg/g dry weight		Male No.	Fe µg/g dry weight	
	Mean*	C.V.**		Mean*	C.V.**
1	167	12.67	26	148	5.15
2	184	12.54	27	183	9.21
3	204	8.25	28	199	25.15
4	201	8.11	29	149	10.28
5	217	9.01	30	162	12.20
6	193	13.73	31	208	9.28
7	205	9.18	32	174	11.11
8	221	9.25	33	172	12.55
9	145	10.21	34	160	6.18
10	188	21.05	35	231	10.14
11	198	20.18	36	153	6.33
12	200	10.11	37	160	14.25

13	182	11.18	38	178	16.6
14	164	9.85	39	192	15.66
15	145	10.85	40	193	15.35
16	222	8.55	41	184	14.20
17	191	9.84	42	205	10.18
18	203	8.95	43	209	10.89
19	175	15.10	44	187	11.33
20	188	12.21	45	161	8.61
21	207	13.15	46	222	14.15
22	202	25.18	47	176	15.04
23	170	14.36	48	206	21.18
24	177	10.15	49	202	10.74
25	216	8.85	50	185	9.78

*Mean of seasons, 2015, 2016 and 2017

**Coefficient of variation

3.1.3. Manganese concentrations

Data presented in Table (3) indicate that pollen grain of date palm male trees contained high concentrations of manganese, the concentration ranging from 97 to 202 $\mu\text{g/g}$ dry weight. The best result obtained from the selective male number (10) which recorded 202 $\mu\text{g/g}$ dry weight respectively as compared to the other male trees.

Table 3. Pollen manganese content of date palm males as the mean of the three seasons

Male No.	Mn $\mu\text{g/g}$ dry weight		Male No.	Mn $\mu\text{g/g}$ dry weight	
	Mean*	C.V.**		Mean*	C.V.**
1	134	15.14	26	129	3.25
2	191	12.10	27	133	5.68
3	182	15.05	28	97	3.54
4	178	12.28	29	136	10.54
5	112	10.18	30	189	10.25
6	98	3.25	31	194	12.25
7	155	5.16	32	203	11.82
8	192	8.65	33	158	9.28
9	141	7.99	34	182	8.92
10	202	12.63	35	179	10.25
11	154	8.15	36	192	9.44
12	177	6.58	37	186	8.15
13	168	9.65	38	172	6.18
14	136	6.85	39	194	10.52
15	121	10.21	40	157	8.25
16	166	13.05	41	142	5.15
17	192	12.15	42	154	4.27
18	143	6.17	43	193	9.88
19	129	5.25	44	171	10.55
20	155	9.18	45	185	11.33
21	181	11.21	46	157	12.25
22	138	10.84	47	142	10.22
23	141	13.13	48	194	10.54
24	174	14.85	49	173	12.12
25	187	17.38	50	184	16.10

*Mean of seasons, 2015, 2016 and 2017

**Coefficient of variation

3.1.4. Copper concentrations

Data presented in Table (4) indicate that pollen grain of date palm male trees contained low concentrations of copper, the concentration ranging from 10 to 30 $\mu\text{g/g}$ dry weight. The highest value which obtained from the selective male number (46) recorded 30 $\mu\text{g/g}$ dry weight respectively as compared to the other male trees.

Table 4. Pollen Copper content of date palm males as the mean of the two seasons

Male No.	Cu $\mu\text{g/g}$ dry weight		Male No.	Cu $\mu\text{g/g}$ dry weight	
	Mean*	C.V.**		Mean*	C.V.**
1	16	3.12	26	14	1.14
2	10	2.16	27	13	1.35
3	22	5.24	28	24	1.58
4	14	2.01	29	21	1.87
5	25	2.18	30	19	1.65
6	32	3.24	31	23	1.24
7	30	3.58	32	15	1.65
8	29	2.66	33	21	1.38
9	11	1.23	34	12	1.59
10	17	1.65	35	26	1.34
11	14	1.21	36	22	2.01
12	13	1.68	37	18	2.35
13	15	1.38	38	29	3.00
14	12	1.65	39	15	1.15
15	18	2.12	40	27	2.01
16	15	2.52	41	23	2.05
17	20	1.32	42	20	2.06
18	19	2.58	43	14	1.32
19	23	2.84	44	28	1.89
20	21	2.44	45	24	1.54
21	24	2.86	46	30	2.06
22	13	2.47	47	15	1.11
23	20	2.38	48	29	1.42
24	23	2.65	49	12	1.11
25	30	2.98	50	25	1.68

*Mean of seasons, 2015, 2016 and 2017

**Coefficient of variation

3.2. Macro nutrient

3.2.1. Magnesium concentrations

Data presented in Table (5) indicated that pollen grain of date palm male trees contained high concentrations of magnesium, the concentration ranging from 1320 to 1985 $\mu\text{g/g}$ dry weight.

Table 5. Pollen magnesium content of date palm males as the mean of the three seasons

Male No.	Mg $\mu\text{g/g}$ dry weight		Male No.	Mg $\mu\text{g/g}$ dry weight	
	Mean*	C.V.**		Mean*	C.V.**
1	1330	16.05	26	1840	16.33
2	1670	17.28	27	1940	15.17
3	1450	18.33	28	1450	12.32
4	1940	18.54	29	1320	14.25
5	1680	18.69	30	1520	15.26
6	1130	19.04	31	1830	17.77

7	1980	19.35	32	1220	18.39
8	1320	16.33	33	1340	14.35
9	1960	16.54	34	1530	16.33
10	1340	18.21	35	1820	17.28
11	1460	16.16	36	1470	17.95
12	1970	15.24	37	1960	16.13
13	1950	15.36	38	1780	16.19
14	1350	16.66	39	1660	18.17
15	1930	16.32	40	1430	13.25
16	1720	17.21	41	1940	16.25
17	1880	19.18	42	1350	11.38
18	1930	19.32	43	1230	16.35
19	1490	14.13	44	1680	10.82
20	1360	12.16	45	1920	18.39
21	1500	13.15	46	1985	10.10
22	1830	14.25	47	1840	11.66
23	1610	14.36	48	1670	15.25
24	1450	15.15	49	1450	13.25
25	1720	16.26	50	1940	16.21

*Mean of seasons, 2015, 2016 and 2017

**Coefficient of variation

The best result which obtained from the selective male number (46) recorded 1985 $\mu\text{g/g}$ dry weight respectively as compared to the other male trees.

4. Discussion

It is clear from the current results that the great male pollinators for using in pollinating female date palms (five seedling date palm males number, 2, 10, 30, 41, 47) was characterized by high content of the estimated element (Zinc, Iron, Manganese, Copper and Magnesium). In this respect, [25] pointed that feather date palm pollen is particularly high mineral content. This could be associated with the high mineral content of the soil horizons in dry areas wherever this species grows. But, the capability of the parent plant to accumulate elements within the pollen is additionally associated with the species. Moreover, [19] found that Egyptian pollen grain contain higher quantity of copper 3.196 mg/g with lower quantity of manganese 2.84mg/g, zinc 2.81mg/g and iron 2.41mg/g. The elements Fe, Mg, Mn, Zn and Cu have been classified as essential elements for the plants [29] and human diet [30]. While not copper, iron cannot be properly reborn to its usable [31], whereas the Mg contain within the pollen grains is also reduced risk of stroke [32]. The presence of Mg^{+2} , Cu^{+3} , Fe^{+2} , Mn^{+2} and Zn^{+2} reflects their perform as essential nutrient components, typically as co-factor activators in metal-ligand enzyme complexes [30]. The healthful importance of pollen grain could also be thanks to its content of Zn that is beneficial to manufacture of testosterone, as antioxidant and essential for traditional functioning of the male reproductive system [30,33], in contribution of Mn [34] which also are essential for traditional functioning of central systema nervosum and are an honest anti-oxidant [30]. Workers, rumored that, iron plays a task in syntheses of RNA, reduction of nitrate to ammonia, [35], chlorophyll synthesis [36], nucleic acid metabolism and chemical change and structural roles of Fe^{++} and Fe^{+++} in plant [37]. Also, iron is either a constituent or a cofactor of many antioxidant enzymes, and might acts as a pro-oxidant issue as a result of free or loosely bound Fe- catalyses free radical generation within the presence of reductants and peroxides through the Fenton reaction. In particular, Fe is involved in the Fe-catalysed Haber–Weiss reaction in which trace amounts of Fe^{3+} are reduced by to produce Fe^{2+} which, in turn, reacts with H_2O_2 to form OH (Fenton reaction). As the intrinsic constituent or metal compound, iron is actively concerned in cellular detoxification reactions catalysed by enzyme, phenolic-dependent

peroxidases (non-specific peroxidases, ascorbate peroxidases and metallic element SOD, that scavenge oxide and superoxide, therefore protective the cell from aerophilous injury. As the intrinsic constituent or metal cofactor, iron is actively concerned in cellular detoxification reactions catalysed by enzyme catalase, phenolic-dependent peroxidases (non-specific peroxidases, ascorbate peroxidases and Fe superoxide dismutase, which scavenge hydrogen peroxide and superoxide, therefore protective the cell from aerophilous injury.. In this respect, [38] reported that, iron deficient helianthus plants appears to have an effect on totally different [the various peroxidase isoenzymes to different extents and to induce a secondary aerobic oxidative stress, as indicated by the redoubled levels of H_2O_2 . On the opposite hand, the foremost vital perform of manganese is said to the oxidation-reduction processes[29].It may be used as a cofactor of many enzymes that act as phosphorylated substrates; conjointly, it plays a task in control the amount of plant hormones (auxins) in plant tissues by activating the auxin oxidation system [35]. The essential role of atomic number 30 is said to the synthesis of essential amino acid organic compound and consequently formation of phytohormone i.e. IAA that act as plant hormone particularly in prolonging height of plants [39].metal has a vital role in supermolecule metabolism, supermolecule synthesis, essential amino acid and IAA synthesis, since it activates variety of enzymes for photosynthesis [40, 41]. The essential role of Zn is related to the synthesis of tryptophan amino acid and consequently formation of phytohormone i.e. IAA that act as plant hormone particularly in prolonging height of plants [39]. Zinc has a vital role in carbohydrate metabolism, protein synthesis, essential amino acid and IAA synthesis, since it activates variety of enzymes for photosynthesis [40, 41]. Also, [42] rumored that, the rise in spore grain from Mg, Zn, Mn, iron and atomic number 29 is various components of the plant. The stimulating impact of micronutrients on plant growth is also because of their role in lepton transmission from water to chlorophyll and manufacturing O_2 gas within the chemical change, additionally to their role within the gas metabolism through activated group enzyme catalyst [43]. Also, the favorable impact of Mg, Zn, Fe, Mn and atomic number 29 on yield and its elements in antecedently studied by [8,9] may be attributed to the rise in photosynthetic pigments concentration [37]. Moreover as, catalyst activity, consequently improvement of plant metabolism [44] Also, [42] reported that, the rise in Pollen grain from Mg, Zn, Mn, Fe and Cu also because of the impact of Zn on synthesis of phytohormone (I.A.A.) that promote ontogenesis process and consequently the amounts of mineral elements absorbed and trans placed into the various components of the plant. The stimulating impact of micronutrients on plant growth is also because of their role in electron transmission from water to chlorophyll and manufacturing oxygen gas within the photosynthesis, additionally to their role within the nitrogen metabolism through activated nitrite reductase enzyme [43]. Also, the favorable impact of Mg, Zn, Fe, Mn and Cu on yield and its components in antecedently studied by [8, 9] may be attributed to the rise in photosynthetic pigments concentration [37] moreover as, enzyme activity, consequently improvement of plant metabolism [44].

5. Conclusions

Finally, from the present results, it could be concluded that the good male pollinators for using in pollinating female date palms was characterized by high content from elements (Zinc, Iron, Manganese, Copper and Magnesium). Which reflected greatly on fertility and yield, and consequently improved quality of fruits because these elements participate in the different metabolic processes. Moreover, it could be recommended to fertilizing date palm males trees with micronutrients of (Fe, Zn, Mn, Cu, Mg) for increase vitality of pollen grains of date Palm male trees and greatly increased yield.

References

- [1] Khoshroo SMR (2011) Seed storage protein electrophoretic profiles in some Iranian date palm (*Phoenix dactylifera* L.) cultivars. African Journal of Biotechnology 10:17793-17804. DOI: 10.5897/ajb11.2726
- [2] Baliga MS, Baliga BRV, Kandathil SM, Bhat HP, Vayalil PK (2011) A review of the chemistry and pharmacology of the date fruits (*Phoenix dactylifera* L.). Food Research International 44: 1812-1822. DOI: 10.1016/j.foodres.2010.07.004
- [3] Mohamed Ahmed MVO, Bouna ZEO, Mohamed Lemine FM, Djeh TKO, Mokhtar T, Mohamed Salem AO (2011) Use of multivariate analysis to assess phenotypic diversity of date palm (*Phoenix dactylifera* L.) cultivars. Scientia Horticulturae 127: 367-371. DOI: 10.1016/j.scienta.2010.11.011
- [4] Ghnaim HD, Al-Muhtaseb JA (2006) Effect of Pollen Source on Yield, Quality and Maturity of 'Mejhool' Date Palm. Jordan Journal of Agricultural Sciences 2:8-15.
- [5] Metwaly HAA, Abou-Rekab ZAM, Abd El-Baky A A, El-Bana AA (2009) Evaluation of some seeded date palm trees grown in Fayoum Governorate .B-chemical characteristics. 4 th Conference on Recent Technologies in Agriculture, 701-715.
- [6] Swaed SY (2009) Superficial study of pollen grains of some agricultural varieties of date palm. Basra Journal of Research Date Palm 8:81-93.
- [7] El-Kosary S (2009) Characteristics of four Barhee date strains as affected by pollen source and pollination time. J. of Hort. Sci. & Ornamental Plants 1:79-91.
- [8] Moustafa AA, Ibrahim ZA, Seif El- Yazel SA, El-Anwar MA (2010a) Evaluation and selection of some seedling Date palm Males Grown at Fayoum Governorate. Proc. 4th int. Date palm Conference Eds. A. Zaid and G.A. Alhadrami Acta Hort. 882.ISHS 2010.
- [9] Moustafa AA, Ibrahim ZA, Seif El-Yazal SA, El- Anwar MA (2010b) Evaluation of different selected seedling date palm males in pollination of "SEEWY" Date palms grown In fayoum Governorate, Egypt .Date palm production and processing technology, pp79-82.2010 Sultan Qaboos University.
- [10] Seif El-Yazal S A, Alharby H F, Seif El-Yazal M A, Hassan G M, Rady M M (2017) Molecular identification of some seedling of date palm (*Phoenix dactylifera* L.) males trees. The Journal of Animal & Plant Sciences, 27:1287-1294.
- [11] Jasim AM, Ibrahim AO, Abbas MF(1995) Certain physicochemical change during growth and maturity of Hillw, date. Basrah. J. Agric. Sci. 8: 2.
- [12] Jasim AM, Yousef AY, Al-Jubouri S (2000) The use of neutron activation analysis technique for estimating protein and mineral elements in the pollen of different varieties of male palm. Basra Agricultural Science Journal 13:41-55.
- [13] Abed MA (2005) Determine of carbohydrates, protein and phenolic compounds content in pollen grains of three of Palm *Phoenix dactylifera*. Basrah Journal For Date Palm Research 4:141-149.
- [14] Abed MA, El-Tememy EH (2006) Determination of three elements in pollen grains of different date palm. Basra Journal of Agricultural Science 19:73-80.
- [15] Abed AM, Abdul Wahid AH, Al-Tamimi EH (2007) Some trace elements concentration (trace) in three varieties of agricultural pollen of date palm. Basra Journal of Agricultural Science 20:7784.

-
- [16] Abed AM, Abdul Wahid AH, Abbas MF (2008) Studing the content of three palm pollen grains varieties from cytokinins, calcium and boron content. Basra Journal of Agricultural Science 2:64-65.
- [17] Abed MA, Hantosh AA, AL-Saad HT, Zadan MA, kames AS (2011) Seasonal variations of some biochemical aspects for five species of date palm (2-trace elements).Basra Science Journal 37:50-66.
- [18] Swaed SY, Aty ME (2011) Determine mineral content for four elements of pollen grains of two date palm cultivars. Basrah Journal For Date Palm Research10 :141-149.
- [19] Hassan HMM (2011) Chemical composition and nutritional value of palm pollen Grains. Global Journal of Biotechnology & Biochemistry 6: 01-07.
- [20] Abdel Rheem DA, Hassan HM, Ahmed SA, Elghandour AH (2015) Investigation and comparison of lipid content and nutritional composition of different organs of *Brahea armata*– S. Watson, Family Arecaceae, growing in Egypt. Journal of American Science 11:87-93. <http://www.jofamericanscience.org>
- [21] Al-Samarai AH, Al-Salihi FG, Al-Samarai RR (2016) Phytochemical constituents and nutrient evaluation of date palm (*Phoenix dactylifera*, L.) pollen grains. Tikrit Journal of Pure Science 21:56-62.
- [22] Talaie A, Badmahmoud AT, Malakout MG (2001) The effect of foliar application of N, B and Zn on quantitative and qualitative characteristics of olive fruit. Iranian J. Agric. Sci. 32: 727-736.
- [23] Wojcik P, Wojcik M (2003) Effect of boron fertilization on conference pear tree vigor, nutrition and fruit yield and storability. Plant and Soil 256: 413-421.
- [24] Aly, H.S.H (2018).Evaluation of pollen grains germination, viability and chemical composition of some date palm males. Middle East J. Agric. Res., 7(2): 235-247.
- [25] Stanley RG, Linskens HF (1974). Pollen biology, chemistry and management, Springerverlage, New York.
- [26] Chapman HD, Pratt PF (1961) Methods of analysis of soil, plants and water. Univ. of California, Riverside, U. S. A. Division of Agricultural Science Berkeley CA.
- [27] Zade MB, Salunke SD(2011) Total mineral content of rare fruits grown in latur district. Hi-Tech Research Analysis 1:6-10.
- [28] Snedecor GW, Cochran WG (1980) Statistical methods. 7th Ed., Iowa State Univ. press, Ames, Iowa, USA.
- [29] 29. Mengel K, Kirkby EA (1982) "Principles of plant narration", 3rd Publisher, International Potash Institute, Worbalaufen- Bern /Switzerland, pp. 593 - 655.
- [30] Zafar M, Khan MA, Ahmad M (2010) Elemental analysis of some medicinal plants used in traditional medicine by atomic absorption spectrophotometer (AAS). Journal of Medicinal Plants Research. 19: 1987-1990.
- [31] Lopez MA, Martos FC(2004) Iron availability: An updated review, International Journal of Food Sciences and Nutrition 55: 597.
- [32] Larsson SC, Orsini N, Wolk A (2012) Dietary magnesium intake and risk of stroke: a meta-analysis of prospective studies. Am J Clin Nutr. 95:362-366.
- [33] Ali H, Ahmed M, Baig M, (2007) Relationship of zinc concentrations in serum and seminal plasma with various semen parameters in infertile subjects. Pak. J. Med. Sci. 23:111-114.

-
- [34] Bansal AK, Kaur ARJ (2009) Cooperative functions of manganese and thiol redox system against oxidative stress in human spermatozoa. J Hum Reprod Sci. 2: 76–80.
- [35] Russell SEW (1989) "Soil conditions and plant growth", 11th Ed. Department of Soil Science, Univ. of Reading, Longman Group, UK Ltd
- [36] Romheld W, Maschnur H (1961) "USA Micronutrients in Agriculture. "2nd Ed. SSSA No. 4. Soil Sci., Society of America, 297.
- [37] Price CA, Clark HE, Funkhouser EA (1972) Function of micronutrients in plants. Soil Sci. Society of America, 231: 136-148.
- [38] Ranieri A, Castagna A, Baldan B, Soldatini G F (2001) Iron deficiency differently affects peroxidase isoforms in sunflower. Journal of Experimental Botany 52:25-35.
- [39] Devendra T, Najda CL, Tak D (1999) Effect of growth regulators on growth and flower yield of tuberose (*Polianthes tuberosa* L.) cv. single. Scientific Hort. 6: 147-150.
- [40] Gardner FP, Pearce RB, Mitchell RL (1985) Physiology of crop plants. The Iowa State Univ. Press., Ames., Iowa, U.S.A.
- [41] Marschner H (1986) Mineral nutrition of higher plants. Academic Press Inc. (London) LTD, pp. 279-287.
- [42] Devlin RM, Withman FH (1985) Plant Physiology. 4th Ed. CBS Publishers and Distribution, 485, Jain Bhawan, Bhola Nath Nagar, Shahdara. Delhi-110 032 (India), p. 443.
- [43] Baza MS (1984) Effect of some macro and micro elements on growth and yields of maize. M.Sc. Thesis, Fac. Agric. of Moshtohr, Zagazig Univ., Egypt.
- [44] Boardman NK (1975) Trace-elements in photosynthesis. In trace elements in Soil- Plant Animal System. Edited by Nicholas: 199-212 Academic Press. Inc., New York . San Francisco and London.